

An Anisotropic Density Turbulence Model from the Sun to 1 au Derived From Radio Observations

EDUARD P. KONTAR,¹ A. GORDON EMSLIE,² DANIEL L. CLARKSON,¹ XINGYAO CHEN,¹ NICOLINA CHRYSAPHI,^{3, 1}
FRANCESCO AZZOLLINI,¹ NATASHA L.S. JEFFREY,⁴ AND MYKOLA GORDOVSKYY⁵

¹*School of Physics & Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK*

²*Department of Physics & Astronomy, Western Kentucky University, Bowling Green, KY 42101, USA*

³*Sorbonne Université, École Polytechnique, Institut Polytechnique de Paris, CNRS, Laboratoire de Physique des Plasmas (LPP), 4 Place Jussieu, 75005 Paris, France*

⁴*Department of Mathematics, Northumbria University, Physics and Electrical Engineering, Newcastle upon Tyne, NE1 8ST, UK*

⁵*Department of Physics, Astronomy & Mathematics. University of Hertfordshire, Hatfield AL10 9AB, UK*

ABSTRACT

Solar radio bursts are strongly affected by radio-wave scattering on density inhomogeneities, changing their observed time characteristics, sizes, and positions. The same turbulence causes angular broadening and scintillation of galactic and extra-galactic compact radio sources observed through the solar atmosphere. Using large-scale simulations of radio-wave transport, the characteristics of anisotropic density turbulence from $0.1 R_{\odot}$ to 1 au are explored. For the first time, a profile of heliospheric density fluctuations is deduced that accounts for the properties of extra-solar radio sources, solar radio bursts, and in-situ density fluctuation measurements in the solar wind at 1 au. The radial profile of the spectrum-weighted mean wavenumber of density fluctuations (a quantity proportional to the scattering rate of radio-waves) is found to have a broad maximum at around $(4-7) R_{\odot}$, where the slow solar wind becomes supersonic. The level of density fluctuations at the inner scale (which is consistent with the proton resonance scale) decreases with heliocentric distance as $\langle \delta n_i^2 \rangle(r) \simeq 2 \times 10^7 (r/R_{\odot} - 1)^{-3.7} \text{ cm}^{-6}$. Due to scattering, the apparent positions of solar burst sources observed at frequencies between 0.1 and 300 MHz are computed to be essentially cospatial and to have comparable sizes, for both fundamental and harmonic emission. Anisotropic scattering is found to account for the shortest solar radio burst decay times observed, and the required wavenumber anisotropy is $q_{\parallel}/q_{\perp} = 0.25 - 0.4$, depending on whether fundamental or harmonic emission is involved. The deduced radio-wave scattering rate paves the way to quantify intrinsic solar radio burst characteristics.

Keywords: interplanetary scintillation (828), interplanetary turbulence (830), radio bursts (1339), solar corona (1483), solar wind (1534)

1. INTRODUCTION

Radio-wave scattering affects the temporal characteristics, sizes, and positions of both solar radio bursts and extra-solar sources observed through the turbulent solar atmosphere. Solar radio bursts (such as Type I, Type II, Type III, etc.) are produced predominantly via plasma mechanisms at frequencies that are close to either the local plasma frequency or its double (harmonic), and are thus particularly strongly affected by scattering (e.g., Fokker 1965; Steinberg 1972; Riddle 1974; Bastian 1994; Arzner & Magun 1999; Thejappa & MacDowall 2008; Kontar et al. 2017; Chrysaphi et al. 2018; Krupar et al. 2018; McCauley et al. 2018; Gordovskyy et al. 2019; Kontar et al. 2019; Chrysaphi et al. 2020; Krupar et al. 2020; Murphy et al. 2021; Maguire et al. 2021; Mohan 2021; Musset et al. 2021; Sharma & Oberoi 2021). Extra-solar point sources observed through the solar atmosphere and solar wind are also noticeably broadened (e.g., Machin & Smith 1952; Hewish 1958; Dennison & Blesing 1972; Anantharamaiah et al. 1994; Sasikumar Raja et al. 2017) and scintillate (Hewish et al. 1964; Cohen et al. 1967; Woo 1977; Coles 1978; Armand et al. 1987; Manoharan 1993a; Breen et al. 1999; Miyamoto et al. 2014; Sasikumar Raja et al. 2016; Chhetri et al. 2018; Tyul’bashev et al. 2023). Overall, various observations provide a qualitatively consistent picture of turbulence in the heliosphere. These observations also strongly suggest that the density fluctuations are anisotropic, with parallel wavenumbers q_{\parallel} that are smaller than the perpendicular wavenumbers q_{\perp} (e.g., Dennison & Blesing 1972; Coles & Harmon 1989; Armstrong et al. 1990), with stronger anisotropy sug-

gested at smaller scales. Observations of solar radio bursts at higher frequencies ($\sim 30\text{--}50$ MHz) originate closer to the Sun and usually require an anisotropy $q_{\parallel}/q_{\perp} \simeq 0.15 - 0.3$ (Kontar et al. 2019; Chen et al. 2020; Kuznetsov et al. 2020; Clarkson et al. 2021), while simultaneous observations of solar radio bursts in interplanetary space using multiple spacecraft require a somewhat larger ratio $q_{\parallel}/q_{\perp} \simeq 0.3 - 0.4$ (Musset et al. 2021). In-situ observations of the wavenumber anisotropy of density fluctuations are very few: Celnikier et al. (1987) report an anisotropy factor between 1 and 2 with an uncertainty comparable to the measured values, while Roberts et al. (2018) report an anisotropy factor between 1 and 3.

The scattering regimes for solar radio bursts and extra-solar point sources are very different. Extra-solar sources are typically observed at frequencies ω that are much greater than the plasma frequency of the solar medium in which their emitted radiation propagates, i.e., $\omega/\omega_{pe} \gg 1$, where $\omega_{pe} = \sqrt{4\pi e^2 n/m_e}$ (s^{-1}) is the local plasma frequency, n (cm^{-3}) is the local number density, and e and m_e are the electronic charge (esu) and mass (g), respectively. The scattering mean free path λ_{sc} is proportional to $(\omega^2 - \omega_{pe}^2)^{-2}$ and is typically much larger than the distance r traveled by the radio waves in the turbulent medium. Consequently, such sources experience weak scattering and their broadening is normally computed using a thin-screen approximation to deduce a wave structure function that measures the spatial correlation of the turbulence at different scales (see, e.g., Lee & Jokipii 1975; Woo et al. 1977; Coles & Harmon 1989). By contrast, plasma emission of solar radio bursts is at frequencies that are inherently close to the local plasma frequency (or its second harmonic), so that $\omega/\omega_{pe} \simeq 1 - 2$. For values of ω so close to ω_{pe} , the mean free path λ_{sc} can be much smaller than the distance a wave propagates between the emission location and the observer, so that the emitted radio waves are (at least close to their origin) subject to multiple scatterings that randomize the wave propagation (e.g., Kontar et al. 2019). Waves emitted in solar radio bursts, at least close to the emitting region, thus propagate diffusively (Bian et al. 2019; Kontar et al. 2019; Chen et al. 2020), with the transport modeled using a stochastic random-phase description (Chandrasekhar 1952; Fokker 1965; Steinberg et al. 1971; Krupar et al. 2018; Thejappa & MacDowall 2008) with inclusion of anisotropic effects (Arzner & Magun 1999; Kontar et al. 2019).

Regarding observations of the density fluctuations, solar corona sounding experiments normally work well for heliocentric distances $\sim (2 - 100) R_{\odot}$ (e.g., Coles & Harmon 1989), but become progressively more difficult at smaller heliocentric distances due to the strong radio emission of the quiet solar corona between $(1 - 2) R_{\odot}$. On the other hand, solar radio burst emissions originating between $(1 - 2) R_{\odot}$ have brightness temperatures far greater than the corona and are routinely observed. As such, solar radio bursts observed over wide range of distances from the Sun, such as Type III bursts, are better suited to provide constraints on scattering and density fluctuations close to the Sun, i.e., below $2 R_{\odot}$ (Kontar et al. 2017; Chrysaphi et al. 2018; Chen et al. 2020; Sharma & Oberoi 2021; Mohan 2021).

A common density turbulence model must self-consistently explain all three sets of observations: (1) extra-solar sources, (2) solar sources, and (3) density fluctuations measured in-situ in the solar wind. Here, using the numerical model of Kontar et al. (2019), we carry out a number of radio-wave propagation simulations between $0.15 - 300$ MHz and we consider the results in light of the very substantial array of extra-solar, solar, and in-situ observations published in the literature. We show that the level of radio-wave scattering necessary to account for the properties of non-solar sources matches the level required to account for the source sizes and apparent positions of solar radio bursts in the corona and in the heliosphere. Further, the burst decay times and time durations predicted by such a turbulence model are consistent with typical Type III burst durations below 1 MHz, with the durations of shortest radio bursts observed (such as spikes and Type IIIb striae) above 1 MHz, and with the Type III burst striae observed by Parker Solar Probe (PSP; Fox et al. 2016), which show a transition from a decay time that is proportional to $1/f$ at lower frequencies to a downshifted $1/f$ behavior at higher frequencies.

In Section 2 we discuss the model of radio-wave propagation in a turbulent medium with an anisotropic turbulence wavenumber spectrum $S(\vec{q})$. In Section 3, we discuss the angular broadening of extra-solar sources with ray paths that pass close to the turbulent atmosphere of the Sun, and we determine the density and turbulence profiles required to account for observations of extra-solar source extents. In Section 4 we discuss simulations of solar radio bursts that use the turbulence profile (both strength and anisotropy) deduced from the data on extra-solar source angular broadening, and we compare the model predictions with observations of source decay times (Section 4.1), sizes (Section 4.2), and apparent positions (Section 4.3). In Section 5 we discuss the relation of the turbulence wavenumber spectrum to in-situ measurements of the frequency spectrum of density fluctuations, leading to further constraints on both the level of turbulence and its degree of anisotropy. In Section 6 we use observations of the turbulence level at the inner scale (the boundary between the inertial and dissipative ranges of the turbulence spectrum) to further constrain the turbulence amplitude, and we present results on the inferred variation of the level of density fluctuations with solar

distance. Finally, in Section 7 we summarize the results obtained and use them to present a self-consistent model of the interplanetary turbulence profile between the Sun and the Earth.

2. RADIO-WAVE TRANSPORT IN AN ANISOTROPIC TURBULENT MEDIUM

Density fluctuations in general are characterized by their three-dimensional wavenumber spectrum $S(\vec{\mathbf{q}})$, such that the density fluctuation variance $\langle \delta n^2 \rangle$, normalized by the local density, is

$$\epsilon^2 \equiv \frac{\langle \delta n^2 \rangle}{n^2} = \int S(\vec{\mathbf{q}}) \frac{d^3 q}{(2\pi)^3}, \quad (1)$$

where $\vec{\mathbf{q}}$ is the wavevector associated with a density fluctuation, $\langle \dots \rangle$ denotes an ensemble average, and $n(\vec{\mathbf{r}})$ is the background plasma density. A typical wavenumber spectrum (e.g., Alexandrova et al. 2013) has three main domains:

$$S(q) = S(q_i) \times \begin{cases} \left(\frac{q}{q_o}\right)^{-3} \left(\frac{q_o}{q_i}\right)^{-11/3}, & q < q_o \quad (\text{outer}) \\ \left(\frac{q}{q_i}\right)^{-11/3}, & q_o \leq q \leq q_i \quad (\text{inertial}) \\ \left(\frac{q}{q_i}\right)^{-2-\delta}, & q > q_i \quad (\text{dissipative}), \end{cases} \quad (2)$$

where the spectrum is normalized to the value $S(q_i)$ at the inner scale wavenumber $q = q_i$, the boundary between the inertial and dissipative ranges. We note that the density fluctuation power spectrum shows evidence of flattening between the inertial and dissipation scales (e.g., Celnikier et al. 1987; Coles & Harmon 1989; Šafránková et al. 2015), but for simplicity we do not consider this here.

The wavenumber volume element $d^3 q \propto q^2 dq$, and, qualitatively, the wavenumber spectrum has three major parts (left panel of Figure 1): $q^2 S(q) \sim 1/q$ for large wavelengths ($q < q_o$), the (Kolmogorov) inertial range $q^2 S(q) \sim q^{-5/3}$ between the outer scale q_o and the inner scale q_i (e.g., Marsch & Tu 1990), and the dissipation range $q^2 S(q) \sim q^{-\delta}$ at small wavelengths ($q > q_i$) (e.g., Celnikier et al. 1983). We will be interested in the spectrum-weighted mean wavenumber $\overline{q\epsilon^2}$ (see Kontar et al. 2019), which is proportional to $\int q S(q) q^2 dq$. This integral can be written as $\int q^4 S(q) d \ln q$; the corresponding integrand is shown schematically in the right panel of Figure 1 and is seen to peak near q_i , corresponding to the inner scale. Quantitatively, the contribution to $\overline{q\epsilon^2} \propto \int_0^\infty q^3 S(q) dq$ from the long-wavelength outer range ($q < q_o$) is $S(q_i) q_i^{11/3} q_o^{-2/3} \int_0^{q_o} dq = S(q_i) q_i^4 (q_o/q_i)^{1/3}$, the contribution from the inertial range ($q_o < q < q_i$) is $S(q_i) q_i^{11/3} \int_{q_o}^{q_i} q^{-2/3} dq = 3 S(q_i) q_i^4 (1 - [q_o/q_i]^{1/3})$, and the contribution from the dissipative range ($q > q_i$) is $S(q_i) q_i^{\delta+2} \int_{q_i}^\infty q^{1-\delta} dq = S(q_i) q_i^4 / (\delta - 2)$. The inertial range between q_o and q_i is typically about three orders of magnitude (e.g., Alexandrova et al. 2013), and a typical dissipation range spectral index δ varies from 2.3 to 2.9 (e.g., Celnikier et al. 1987), so we adopt $\delta = 2.5$. The contributions to $\overline{q\epsilon^2}$ from the three ranges are therefore in the approximate ratio 0.1 : 3 : 2. Ignoring the small contribution at large scales ($q < q_o$), the spectrum-weighted mean wavenumber $\overline{q\epsilon^2}$ becomes

$$\overline{q\epsilon^2} = \int S(q) \frac{d^3 q}{(2\pi)^3} \simeq \frac{4\pi}{(2\pi)^3} \left(3 + \frac{1}{\delta - 2}\right) S(q_i) q_i^4 = \frac{20\pi}{(2\pi)^3} S(q_i) q_i^4, \quad (3)$$

where the last equality uses $\delta = 2.5$, as discussed above. Equation (3) and Figure 1 importantly highlight that $\overline{q\epsilon^2}$ is determined mostly by density fluctuations close to the inner scale q_i^{-1} .

Observations often suggest the presence of anisotropy relative to the direction of the magnetic field, suggesting a correspondingly anisotropic wavenumber spectrum. Following previous studies (Dennison & Blesing 1972; Woo et al. 1977), we therefore consider a wavenumber spectrum of the form

$$S(\vec{\mathbf{q}}) = S(\tilde{q}), \quad \text{where} \quad \tilde{q} = \sqrt{q_{\perp 1}^2 + q_{\perp 2}^2 + \frac{q_{\parallel}^2}{\alpha^2}}, \quad (4)$$

which has axial symmetry around the \parallel direction, i.e., along the (assumed radial) magnetic field $\vec{\mathbf{B}}$. The wavenumber diffusion tensor describing elastic scattering of radio waves with wavenumber $\vec{\mathbf{k}}$ in such an anisotropic medium can be written (e.g., Arzner & Magun 1999; Bian et al. 2019; Kontar et al. 2019)




Figure 1. *Left:* Spectrum of density fluctuations showing the $q^2 S(q) \sim q^{-1}$ spectrum at large scales $< q_o^{-1}$, the (Kolmogorov) inertial range $q^2 S(q) \sim q^{-5/3}$ between the outer scale q_o^{-1} and the inner scale q_i^{-1} , and the dissipation range $q^2 S(q) \sim q^{-\delta}$ (with $\delta > 2$) at small scales $> q_i^{-1}$. *Right:* The $\ln q$ space integrand required to compute the mean wavenumber $\overline{q\epsilon^2} \propto \int q S(q) q^2 dq = \int q^4 S(q) d \ln q$, highlighting the peak near the inner scale q_i^{-1} . In both plots, both axes are on logarithmic scales.

$$D_{ij} = \frac{\omega_{pe}^4}{32 \pi^2 \omega c^2} \int q_i q_j S(\vec{q}) \delta(\vec{q} \cdot \vec{k}) d^3 q, \quad (5)$$

where $\omega(\vec{k}) = \sqrt{\omega_{pe}^2 + c^2 k^2}$ is the angular frequency and \vec{k} the wavevector of electromagnetic waves in a plasma with local plasma frequency $\omega_{pe}(r)$. For the spectrum given by Equation (4), we can obtain (Kontar et al. 2019) an explicit expression for the wavenumber diffusion tensor components:

$$D_{ij} = \left[\frac{A_{ij}^{-2}}{(\vec{k} \mathbf{A}^{-2} \vec{k})^{1/2}} - \frac{(\mathbf{A}^{-2} \vec{k})_i (\mathbf{A}^{-2} \vec{k})_j}{(\vec{k} \mathbf{A}^{-2} \vec{k})^{3/2}} \right] \frac{\pi \omega_{pe}^4}{16 \omega c^2} \alpha \int \tilde{q} S(\tilde{q}) \frac{d^3 \tilde{q}}{(2\pi)^3}, \quad (6)$$

where $\tilde{\mathbf{q}} = (q_{\perp,1}, q_{\perp,2}, q_{\parallel}/\alpha)$ is the wavevector transformed into a space such that $S(\tilde{q})$ is isotropic, \mathbf{A} is the matrix $\text{diag}(1, 1, \alpha^{-1})$, and we have used $d^3 q = \alpha d^3 \tilde{q}$. The case $\alpha = 1$ corresponds to isotropic turbulence, while the typically observed case $\alpha < 1$ has power predominantly oriented in the perpendicular direction. In such a situation, the density perturbations are elongated along the field lines, so that the parallel length is larger than the perpendicular length: $q_{\parallel}^{-1} > q_{\perp}^{-1}$, i.e., the perpendicular wavenumber is larger than the parallel wavenumber: $q_{\parallel} < q_{\perp}$.

3. ANGULAR BROADENING OF EXTRA-SOLAR RADIO SOURCES

We consider the propagation of a radio wave from a distant extra-solar source with a heliocentric angular separation of $\simeq 0.27 r/R_{\odot}$ degrees, where r is the linear distance of closest approach of the ray path to the center of the Sun and 0.27° is the angular radius of the Sun (see Figure 2). Following Equation (4), we define the q_{\parallel} direction as that aligned with the (assumed radial) solar magnetic field $\vec{\mathbf{B}}$, and we further define two directions perpendicular to this axial direction: $q_{\perp,2}$ in the direction \hat{y} on the plane of the sky that is perpendicular to $\vec{\mathbf{B}}$ (Figure 2), and $q_{\perp,1}$ in a direction orthogonal to both q_{\parallel} and $q_{\perp,2}$ (which, at the point of closest approach, is in the same direction as the wave propagation direction \hat{z}). Evidently, radio waves propagating from the source will be affected by scattering in the turbulent solar atmosphere and, for emission frequencies ω that are much larger than the plasma frequencies ω_{pe} along the ray path though the heliosphere, this can be considered in the weak-scattering limit. In this limit, Equation (5) shows that the angular broadening rates per unit travel distance $dz = v_{gr} dt = (c^2 k/\omega) dt$, in the direction¹ \perp_2 (perpendicular to the radial direction on the plane of the sky), can be written, using Equation (6) (see also Kontar et al. 2019),

$$\frac{d\langle \theta_{\perp,1}^2 \rangle}{dz} = \frac{2 D_{\perp,1}}{v_{gr} k^2} = \frac{\pi}{8} \frac{\omega_{pe}^4}{v_{gr} c^2 \omega k^3} \frac{1}{(2\pi)^3} \int q_{\perp,2}^2 S(q_{\perp,1} = 0, q_{\perp,2}, q_{\parallel}) dq_{\parallel} dq_{\perp,2} = \frac{\pi}{8} \frac{\omega_{pe}^4}{\omega^4} \overline{q \epsilon^2}, \quad (7)$$

¹ The angular broadening rate in the parallel (radial) direction on the plane of the sky can be evaluated similarly by integrating the projections of $D_{\perp,1}$ and $D_{\parallel\parallel}$ along the line of sight.

where $\vec{\nabla}_{gr} = \partial\omega/\partial\vec{\mathbf{k}} = c^2 \vec{\mathbf{k}}/\omega$ is the group velocity of the radio wave and we have used the approximate (high-frequency, i.e. $\omega \gg \omega_{pe}$) dispersion relation $\omega = ck$. For a spherically symmetric corona, the apparent rms source sizes in the perpendicular direction, observed at frequency $f = \omega/2\pi$ (Hz), thus satisfy

$$\langle\theta_{\perp}^2\rangle^{1/2}(r) \propto \frac{1}{f^2} \left[\int_r^{\infty} \overline{q\epsilon^2}(r') n^2(r') dr' \right]^{1/2}, \quad (8)$$

where the constant of proportionality depends² on the distance of closest approach r of the ray to the Sun, and we have used the relation $\omega_{pe} \propto n^{1/2}$. At radial distance r , the mean (spectrum-averaged) wavenumber, $\overline{q\epsilon^2}$, at the position r is given by substituting Equation (4) into Equation (1):

$$\overline{q\epsilon^2} R_{\odot} = R_{\odot} \int S(q) \frac{d^3q}{(2\pi)^3} = \alpha R_{\odot} \int \tilde{q} \mathcal{S}(\tilde{q}) \frac{d^3\tilde{q}}{(2\pi)^3} = \alpha \frac{4\pi}{(2\pi)^3} R_{\odot} \int_0^{\infty} \tilde{q}^3 S(\tilde{q}) d\tilde{q}, \quad (9)$$

where we have multiplied $\overline{q\epsilon^2}$ by the solar radius R_{\odot} to produce a dimensionless quantity $\overline{q\epsilon^2} R_{\odot}$. The angular broadening rates both parallel and perpendicular to the magnetic field are thus both proportional to the square root of the quantity $\overline{q\epsilon^2} R_{\odot}$.




Figure 2. *Left:* Schematic showing the coordinate system used and its relation to the solar disk and the line of sight to an extra-solar point source with separation distance r . The broadening of point sources is calculated as an integral along z in Appendix B. *Right:* Observed FWHM major-axis source sizes of broadened point sources (Hewish 1958; Slee 1959; Högbom 1960; Gorgolewski et al. 1962; Hewish & Wyndham 1963; Slee 1966; Harries et al. 1970; Blesing & Dennison 1972; Woo 1978; Bradford & Routledge 1980; Narayan et al. 1989; Anantharamaiah et al. 1994). The Hewish (1958) data shows points where the detector baseline orientation was specified, and we take the larger of the measurements as the assumed major axis dimension. The solid black line shows the data fit $\theta = 355(r/R_{\odot} - 1)^{-2.15}$ arcsec, and extrapolated by the dashed line. The colored lines show the simulated data for various values of the anisotropy parameter α , with the lower and upper bounds of each region denoting values of $\overline{q\epsilon^2} R_{\odot}$ corresponding to $0.5\times$ and $2\times$ the nominal values from Equation (10). Different observations are scaled to a frequency of 1.5 GHz using a frequency dependence f^{-2} , as expected from Equation (7); see also Equation (B8) in Appendix B.

For a prescribed wave frequency f and distance of closest approach r , Equation (8) shows that the source rms angular size and shape depend only on the profiles of plasma density n (through the local plasma frequency ω_{pe})

² See Appendix B for a more formal expression for Equation (8).

and the spectrum-weighted mean wavenumber of density fluctuations $\overline{q\epsilon^2} R_\odot$, at distances larger than the closest ray-approach distance. The turbulence profile in turn depends on the spectrum $S(\tilde{q})$ and its degree of anisotropy, characterized by the parameter α (see Equation (4)). Further, Equation (8) shows that the apparent angular extent of the source in the radial direction is smaller than the apparent extent in the perpendicular direction on the plane of the sky, by a factor α . Hereafter we define the source “size” as the major axis, i.e., the FWHM source extent in the y -direction, perpendicular to the radial direction on the plane of sky.

If the density and turbulence profiles can be reasonably represented by power-law forms ($n(r) \sim r^{-\gamma}$ and $\overline{q\epsilon^2}(r) \sim r^{-\zeta}$, respectively), then Equation (8) shows that $\langle\theta^2\rangle^{1/2} \sim r^{-[\gamma+(\zeta-1)/2]}$. The data on the apparent sizes of extra-solar sources are presented in Figure 2; they show that $\langle\theta^2\rangle^{1/2} \propto r^{-2.15}$ for $r \gg R_\odot$. Thus, with a radial density profile $n(r) \propto r^{-2.3}$ (valid at $r \gg R_\odot$; see Appendix A), Equation (8) requires that $(\zeta - 1)/2 = 2.15 - 2.3 = -0.15$; i.e., $\zeta = 0.7$. We therefore adopt the nominal turbulence profile

$$\overline{q\epsilon^2} R_\odot = 2 \times 10^3 \alpha \left(\frac{R_\odot}{r}\right)^{0.7} \left(1 - \frac{R_\odot}{r}\right)^{2.7}, \quad (10)$$

which is proportional to α (Equation (9)) and also includes an additional factor so that $\overline{q\epsilon^2}$ decreases towards the solar surface. This latter factor is required by the solar radio burst observations discussed in the next section. The scaling factor 2×10^3 and the exponent 2.7 in the additional term are chosen to best match the simulation results with the observational data in Figure 2.

4. SOLAR RADIO BURSTS

Radio waves emitted by solar sources (e.g., Type III bursts) suffer scattering in the same turbulent environment that leads to the angular broadening of extra-solar sources. Because such bursts are produced by plasma processes at or near the plasma frequency ω_{pe} or its double (see, e.g., Ginzburg & Zhelezniakov 1958), the scattering process is much stronger than for extra-solar sources with $\omega \gg \omega_{pe}$ and hence the weak-scattering treatment of the previous section is not applicable. Here, using a code developed by Kontar et al. (2019), we numerically simulate the propagation of radio waves in the presence of a prescribed density fluctuation profile $\overline{q\epsilon^2}$ as function of r and anisotropy parameter α , which we take³ to be everywhere constant in our model. We performed simulations with $\overline{q\epsilon^2} R_\odot$ scaled from its nominal value in Equation (10) by a factor in the range $[1/4, 4]$, and using anisotropy parameters α in the range $[0.19, 0.42]$. The results of these simulations lead to predictions of solar radio source properties (source size, apparent position, time profile) as seen from 1 au, and are compared with observations to ascertain the veracity of the $\overline{q\epsilon^2} R_\odot$ profile and anisotropy parameter α used. For the simulations, we assume an initially isotropic distribution of photons emitted near the plasma frequency (fundamental), or twice the plasma frequency (harmonic). Assuming a spherically symmetric corona, we ran numerical simulations of the wave propagation, taking into account anisotropic scattering, large scale refraction, and free-free absorption, using profiles for the ambient density profile $n(r)$ given in Appendix A. We performed simulations for radio frequencies covering the range from ~ 0.15 MHz to ~ 300 MHz, thus encompassing ground-based and space-based (below the ionospheric cut-off) observations. We simulated burst decay times (Subsection 4.1), angular source sizes (Subsection 4.2), and source positions (Subsection 4.3), which are then compared with pertinent observations to determine the spectrum-weighted mean wavenumber of density fluctuations $\overline{q\epsilon^2} R_\odot$ as a function of r , and anisotropy parameter α , that together best account for the observations.

4.1. Decay Time

The top panels of Figure 3 show predicted $1/e$ decay times for the range of simulations discussed above, both for sources that emit at the fundamental (left panel) and for those that emit at the second harmonic (right panel).

The decay time of solar radio bursts has been observed both from space and from the ground, resulting in an extended set of decay-time observations. The bottom panels of Figure 3 compare observations of characteristic decay times (expanding the collection of decay times presented by Kontar et al. 2019) with those obtained from our simulations, assuming emission either at the fundamental (left panel) or at the harmonic (right panel). The vertical bars in Figure 3 show the spread of measurements (where a spread is present), rather than measurement uncertainties. The decay profile is normally well-fit by an exponential form, with a generally well-measured characteristic decay time (e.g., Krupar et al. 2018, 2020) both below 1 MHz and above 20 MHz, with a gap in near-Earth data between a few MHz

³ In principle, α could be treated as a function of r in the simulations, but this would make the parameter space too large to study effectively.

(due to the limited time-resolution of space-based observations) and $\sim 10 - 20$ MHz (due to the ionospheric cut-off for ground-based observations). Although there is a noticeable spread in the observations, the grey area, representing a factor of $0.5\times$ to $2\times$ the nominal $\overline{q\epsilon^2} R_\odot$ profile of Equation (10), includes all data below 3 MHz and is found close to the observed decay times of the high-frequency spikes and Type IIIb fine structures. This is convincing evidence that scattering effects account for the shortest observed decay times. The decay time of an average Type III burst at > 10 MHz is normally quite a bit larger, and is principally determined by the beam width and/or emission processes (e.g., Zhang et al. 2019; Chen et al. 2020). Figure 3 suggests that if all observations are due to harmonic emission, the simulations require $\alpha \simeq 0.4$, while the assumption of emission at the fundamental requires a smaller anisotropy $\alpha \simeq 0.25$ to explain decay time observational data.

4.2. Angular Size

The top panels of Figure 4 show predicted apparent source sizes for the range of simulations discussed above, both for sources that emit at the fundamental (left panel) and for those that emit at the second harmonic (right panel). We have expanded the collection of Type III solar radio burst angular extents initially presented by Kontar et al. (2019). Source sizes at frequencies above 10 MHz are evaluated using ground-based interferometers, while those below 1 MHz are determined using indirect techniques (e.g., spinning demodulation and/or goniopolarimetric techniques; Alvarez 1976; Bougeret et al. 2008; Cecconi 2014). Similarly to the decay time measurements of Figure 3, the bars in Figure 4 show the spread of measurements (where a spread is present), rather than measurement uncertainties. The bottom panels of Figure 4 show observed FWHM sizes (given as the FWHM semi-major axis as observed at 1 au) compared with the predictions of the simulations, assuming emission either at the fundamental (left panel) and at the harmonic (right panel). The simulated source sizes are given in terms of their FWHM (for sources with emission frequencies below 0.15 MHz, the source is so large that it extends all the way to the observer at 1 au). One can see that the majority of observations are, similar to the decay time observations of Section 4.1, within the range $0.5\times$ to $2\times$ the nominal $\overline{q\epsilon^2} R_\odot$ profile of Equation (10).

Publications of observations of Type III burst decay times and/or source sizes rarely identify whether the observations correspond to emission at the fundamental or at the second harmonic, and we therefore conservatively include the possibility of either in our comparison of observed decay times and sizes with simulation results. With that in mind, it is interesting to note that both fundamental and harmonic sources have similar sizes over a wide range of frequencies. This is because the scattering surfaces that principally determine the source size are located at levels where both the plasma frequency and its double are significantly less than the emission frequency, so that whether the emission is fundamental or harmonic is not a critical factor in determining the apparent source size. This also explains why harmonic and fundamental Type III burst source sizes measured at the same frequency are nearly identical (Dulk & Suzuki 1980; Kontar et al. 2017; Chen et al. 2023).

4.3. Apparent Position

Since the early observations of Type III bursts, it was noted (e.g., Shain & Higgins 1959; Fainberg & Stone 1974) that the source positions are not coincident with the positions in the solar atmosphere and solar wind with densities corresponding to emission at the plasma frequency (or its double); equivalently, the observed burst frequencies required a higher plasma density than that is normally observed by optical telescopes (e.g., Fainberg & Stone 1974). This is because radio wave scattering substantially shifts the burst emission away from the Sun, so that the density inferred by associating the burst emission with the plasma frequency (or its double) could be 4-5 times higher than the density at the apparent burst position (e.g., Chrysaphi et al. 2018). Figure 5 shows the variation of observed source positions versus frequency, compared to the predictions of our simulations under the assumptions that the emission is at the plasma frequency (fundamental mode; left panel) or its second harmonic (right panel). We see that the radio-wave scattering reproduces the observed positions over a wide range of distances from the low corona into the solar wind, as shown by the grey band; only a relatively small spread, spanning the range $0.25\times$ to $4\times$ the nominal $\overline{q\epsilon^2} R_\odot$ profile of Equation (10), is required to explain the available observational data points.

These simulations also highlight that, similar to source sizes (but unlike burst decay times), the simulated apparent source positions are relatively independent of the assumption of whether fundamental or harmonic emission is involved. Given an observation frequency, if there were no scattering, sources corresponding to emission at the harmonic should be further outward than sources corresponding to emission at the fundamental. However, in the presence of strong




Figure 3. *Top panels:* Simulated solar radio burst $1/e$ decay times versus frequency. The simulations were performed for turbulence profiles $q\epsilon^2 R_\odot$ given by Equation (10) multiplied by factors $[1/4, 1/2, 1, 2, 4]$, and values of the anisotropy parameter $\alpha = [0.19, 0.25, 0.33, 0.42]$, and for both fundamental (left panels) and harmonic emission (right panels). *Bottom panels:* Observations of source decay times. The solid black points show measurements of Type III decay times (Alexander et al. 1969; Elgaroy & Lyngstad 1972; Alvarez & Haddock 1973a; Barrow & Achong 1975; Krupar et al. 2018; Reid & Kontar 2018), while open black points show spike decay times (McKim Malville et al. 1967; Barrow et al. 1994; Shevchuk et al. 2016; Clarkson et al. 2023). The solid black lines represent power-law fits to the spike data from Equation (3) of Clarkson et al. (2023) and to the Type III data from Equation (51) of Kontar et al. (2019). The light blue points show Type IIIb striae median decay times observed by Parker Solar Probe between 0.95 and 2.15 MHz. The colored lines represent simulated decay times from anisotropic scattering simulations at the fundamental (left panel) and harmonic (right panel). The lower and upper bound of each region denotes values corresponding to turbulence profiles of $0.5q\epsilon^2 R_\odot$ and $2q\epsilon^2 R_\odot$, respectively.




Figure 4. *Top panels:* Simulated FWHM source sizes of Type III bursts. The simulations were performed for turbulence profiles $\overline{q\epsilon^2} R_\odot$ given by Equation (10), multiplied by factors of [1/4, 1/2, 1, 2, 4], for values of the anisotropy parameter $\alpha = [0.19, 0.25, 0.33, 0.42]$, and for both fundamental (left panels) and harmonic emission (right panels). *Bottom panels:* average FWHM size observations (Bougeret et al. 1970; Abranin et al. 1976; Alvarez 1976; Abranin et al. 1978; Chen & Shawhan 1978; Dulk & Suzuki 1980; Steinberg et al. 1985; Saint-Hilaire et al. 2013; Krupar et al. 2014; Kontar et al. 2017) and spikes (Clarkson et al. 2023). Simulated sizes from anisotropic scattering simulations at the fundamental (left) and harmonic (right) are overlaid with colored lines at various values of α , with turbulence levels ranging from $0.5 \overline{q\epsilon^2} R_\odot$ (lower) to $2 \overline{q\epsilon^2} R_\odot$ (upper), where the nominal $\overline{q\epsilon^2} R_\odot$ profile is given by Equation (10). The grey region shows the spread for $\alpha = 0.25$, that is consistent with the observations.




Figure 5. Observed source position versus frequency (Wild et al. 1959; Shain & Higgins 1959; Morimoto & Kai 1961; Gurnett et al. 1978; Bougeret et al. 1984a; Reiner et al. 1998, 2009; Chrysaphi et al. 2018). The colored lines represent the observed positions from the simulations for the fundamental (left) and harmonic (right) at various values of the anisotropy parameter α and the turbulence profile $\overline{q\epsilon^2} R_\odot$, multiplied by 1/4 (lower bound) to 4 (upper bound). The grey region shows the spread across values for an anisotropy parameter $\alpha = 0.25$. The black dashed and dotted lines show the locations corresponding to the fundamental (plasma frequency) and harmonic (double plasma frequency), respectively.

scattering (Figure 5), the apparent source positions are significantly further away from the Sun than the locations expected for either emission at the harmonic mode or at the fundamental. This effect is noticeable for all except the highest emission frequencies $f > 100$ MHz, where the apparent source position is close to the location expected from emission at the harmonic. This shows clearly that measured source locations are dominated by scattering effects and succinctly explains the observations reported by Stewart (1972), showing that the radial distances for fundamental and harmonic components at 80 MHz are, on average, equal.

4.4. Remarks

The observations of Type III burst decay time, source size and position (Figures 3 through 5) show a noticeable data gap at frequencies between ~ 5 MHz and ~ 15 MHz. Observations below the ionospheric cut-off are made from space, while observations above 10 – 15 MHz are ground-based. Parker Solar Probe (PSP; Fox et al. 2016) has sufficiently high frequency resolution to resolve Type IIIb burst striae (Pulupa et al. 2020). Our analysis of 10 Type IIIb bursts (58 striae from April 2-9, 2019) provides interesting measurements (blue symbols in Figure 3), suggesting that Type IIIb and Type III bursts have nearly identical decay times below 1 MHz, but the fine structures in Type IIIb bursts decay faster at higher frequencies. This suggests that the Type III decay is purely due to scattering below 1 MHz and that electron transport is of relevance at higher frequencies. In other words, the scattering of Type III radio bursts is so large below 1 MHz so that it dominates the decay time.

The profiles of $\overline{q\epsilon^2} R_\odot$ (see Figure 6) necessary to reproduce the observations of the source size, decay time and source position are consistent with an anisotropic scattering model with $\alpha \simeq 0.25$ (assuming emission at the fundamental) and $\alpha \simeq 0.42$ (assuming purely harmonic emission). We therefore conclude that turbulent density fluctuations, with a magnitude profile consistent with Equation (10) and an anisotropy parameter $0.2 \lesssim \alpha \lesssim 0.4$, are present in the solar corona and heliosphere.

5. IN-SITU OBSERVATIONS OF DENSITY FLUCTUATIONS

Density fluctuations are often measured in the solar wind (e.g., Celnikier et al. 1983; Marsch & Tu 1990) and can be compared to our density turbulence model by converting the wavenumber spectrum into the frequency power spectrum




Figure 6. *Left:* Spectrum-averaged wavenumber $\overline{q\epsilon^2} R_\odot$ as a function of heliocentric distance r , as given by Equation (10), for fundamental emission with an anisotropy factor $\alpha = 0.25$ (solid), and for harmonic emission with $\alpha = 0.42$ (dashed). The grey region denotes the range from $[1/2, 2] \times \overline{q\epsilon^2} R_\odot$. The data from Coles & Harmon (1989) are retrieved from the density spectrum in their Figure 4, with a density profile $n(r)$ taken from equation (A1). The light grey points are not well constrained due to a lack of observations at the (dominant; see Section 2) inner scale wavenumber q_i . The in-situ density fluctuations data at 1 au (Celnikier et al. 1983, 1987; Kellogg & Horbury 2005; Chen et al. 2012) are retrieved using Equation (20), with the black points corresponding to $\alpha = 0.25$ (fundamental) and the red points corresponding to $\alpha = 0.42$ (harmonic). The data for Celnikier et al. (1987) are taken from their Figure 7. We use the Kellogg & Horbury (2005) spectra for densities $(5-6) \text{ cm}^{-3}$ and $(10-15) \text{ cm}^{-3}$. *Right:* As in the left panel but multiplied by n^2 , with $n(r)$ given by Equation (A1). The dotted line shows the relation $\overline{q\epsilon^2} R_\odot n^2(r) = 6.5 \times 10^{14} (r/R_\odot - 1)^{-5.17}$.

$P(f)$ ($\text{cm}^{-6} \text{ Hz}^{-1}$) measured by a spacecraft as the solar wind, with velocity \vec{V}_{SW} , advects the turbulent fluctuations past the measuring instrumentation. $P(f)$ is related to the wavenumber spectrum $S(\vec{q})$ of density fluctuations by (e.g., Fredricks & Coroniti 1976):

$$P(f) = n^2 \int S(\vec{q}) \delta\left(f - \frac{\vec{q} \cdot \vec{V}_{\text{SW}}}{2\pi}\right) \frac{d^3q}{(2\pi)^3}. \quad (11)$$

In Appendix C, we evaluate $P(f)$ for several illustrative forms (both isotropic and anisotropic) of $S(\vec{q})$. Here we explore the reverse problem of determining the value of $\overline{q\epsilon^2} R_\odot$ from in-situ observations of $P(f)$. We first transform the integration variable from $\vec{q} = (q_{\perp 1}, q_{\perp 2}, q_{\parallel})$ to $\tilde{\vec{q}} = (q_{\perp 1}, q_{\perp 2}, q_{\parallel}/\alpha)$ and define $\tilde{\vec{V}}_{\text{SW}} = (V_{\text{SW}\perp 1}, V_{\text{SW}\perp 2}, \alpha V_{\text{SW}\parallel})$, so that $\tilde{\vec{q}} \cdot \tilde{\vec{V}}_{\text{SW}} = \vec{q} \cdot \vec{V}_{\text{SW}}$ and $d^3q = \alpha d^3\tilde{q}$. Then, since $S(\tilde{\vec{q}})$ is (by construction⁴) isotropic,

$$P(f) = \frac{\alpha n^2}{(2\pi)^2} \int_{\tilde{q}_{\perp}=0}^{\infty} \int_{\tilde{q}_{\parallel}=-\infty}^{\infty} S\left(\sqrt{\tilde{q}_{\perp}^2 + \tilde{q}_{\parallel}^2}\right) \delta\left(\frac{\tilde{q}_{\parallel} \tilde{V}_{\text{SW}}}{2\pi} - f\right) \tilde{q}_{\perp} d\tilde{q}_{\perp} d\tilde{q}_{\parallel}, \quad (12)$$

this can be written as

$$\begin{aligned} P(f) &= \frac{\alpha n^2}{2\pi \tilde{V}_{\text{SW}}} \int_{\tilde{q}_{\perp}=0}^{\infty} \int_{\tilde{q}_{\parallel}=-\infty}^{\infty} S\left(\sqrt{\tilde{q}_{\perp}^2 + \tilde{q}_{\parallel}^2}\right) \delta\left(\tilde{q}_{\parallel} - \frac{2\pi f}{\tilde{V}_{\text{SW}}}\right) \tilde{q}_{\perp} d\tilde{q}_{\perp} d\tilde{q}_{\parallel} \\ &= \frac{\alpha n^2}{2\pi \tilde{V}_{\text{SW}}} \int_{\tilde{q}_{\perp}=0}^{\infty} S\left(\left[\tilde{q}_{\perp}^2 + \left(\frac{2\pi f}{\tilde{V}_{\text{SW}}}\right)^2\right]^{1/2}\right) \tilde{q}_{\perp} d\tilde{q}_{\perp} \end{aligned}$$

⁴ The analysis of Equations (12) through (18) thus applies for the isotropic spectrum $P(f)$ of Appendix C.1 and even for the anisotropic spectrum of Appendix C.2 (which can be scaled into an isotropic form), but not for the spectrum of Appendix C.3, which cannot be transformed into an isotropic form.

$$= \frac{\alpha n^2}{2\pi \tilde{V}_{\text{SW}}} \int_{\tilde{q}=2\pi f/\tilde{V}_{\text{SW}}}^{\infty} \tilde{q} S(\tilde{q}) d\tilde{q} , \quad (13)$$

where in the last equality we have used the substitution $\tilde{q}^2 = \tilde{q}_\perp^2 + (2\pi f/\tilde{V}_{\text{SW}})^2$.

We can now construct the moments of the frequency spectrum of the observed density fluctuations:

$$\int_0^{\infty} f^m P(f) df = \frac{\alpha n^2}{2\pi \tilde{V}_{\text{SW}}} \int_{f=0}^{\infty} f^m df \int_{\tilde{q}=2\pi f/\tilde{V}_{\text{SW}}}^{\infty} \tilde{q} S(\tilde{q}) d\tilde{q} , \quad (14)$$

and reversing the order of integration gives

$$\int_0^{\infty} f^m P(f) df = \frac{\alpha n^2}{2\pi \tilde{V}_{\text{SW}}} \int_{\tilde{q}=0}^{\infty} \tilde{q} S(\tilde{q}) d\tilde{q} \int_0^{\tilde{q}\tilde{V}_{\text{SW}}/2\pi} f^m df . \quad (15)$$

Two moments are of particular interest. First, for $m = 0$,

$$\int_0^{\infty} P(f) df = \frac{\alpha n^2}{(2\pi)^2} \int_{\tilde{q}=0}^{\infty} \tilde{q}^2 S(\tilde{q}) d\tilde{q} = \alpha \frac{\langle \delta n^2 \rangle}{2} , \quad (16)$$

where we have used Equation (1). The factor of 2 arises because only positive frequencies are considered (e.g., for an isotropic distribution, $\langle \delta n^2 \rangle = 2 \int_0^{\infty} P(f) df = \int_{-\infty}^{\infty} P(f) df$). Second, for $m = 1$,

$$\int_0^{\infty} f P(f) df = \frac{\alpha n^2 \tilde{V}_{\text{SW}}}{2(2\pi)^3} \int_{\tilde{q}=0}^{\infty} \tilde{q}^3 S(\tilde{q}) d\tilde{q} , \quad (17)$$

and comparing this to the expression (9) for $\overline{q \epsilon^2} R_\odot$, we immediately see that

$$\overline{q \epsilon^2} R_\odot = 4 \frac{2\pi}{n^2 \tilde{V}_{\text{SW}}} R_\odot \int_0^{\infty} f P(f) df . \quad (18)$$

Now, $\tilde{V}_{\text{SW}} = V_{\text{SW}} (\sin^2 \theta_B + \alpha^2 \cos^2 \theta_B)^{1/2}$, where θ_B is the angle between the solar wind velocity and the (axis of symmetry) magnetic field. Thus, if $\theta_B = 0$,

$$\overline{q \epsilon^2} R_\odot |_{\theta_B=0} = \frac{4}{\alpha} \frac{2\pi}{n^2 V_{\text{SW}}} R_\odot \int_0^{\infty} f P(f) df , \quad (19)$$

while, for a typical 1 au angle $\theta_B \simeq 45^\circ$,

$$\overline{q \epsilon^2} R_\odot |_{\theta_B=45^\circ} = 4 \sqrt{\frac{2}{1+\alpha^2}} \frac{2\pi}{n^2 V_{\text{SW}}} R_\odot \int_0^{\infty} f P(f) df . \quad (20)$$

The frequency-weighted integral over the measured fluctuation power spectrum $P(f)$ at 1 au can be readily evaluated numerically using the data from the spacecraft measurements of [Celnikier et al. \(1983\)](#), [Celnikier et al. \(1987\)](#), [Kellogg & Horbury \(2005\)](#), and [Chen et al. \(2012\)](#), allowing $\overline{q \epsilon^2} R_\odot$ to be determined at 1 au. The resulting values, for various values of α , have been added to Figure 6, and are consistent with the general form of $\overline{q \epsilon^2} R_\odot$ in Equation (10).

6. AMPLITUDE OF DENSITY FLUCTUATIONS

Here we provide further observational constraints on the profile of the spectrum-weighted mean wavenumber of density fluctuations $\overline{q \epsilon^2} R_\odot$, using both observations of the (dominant – Section 2) inner scale at the boundary between the inertial and dissipative ranges of the turbulence wavenumber spectrum (Equation (2)), and published results related to scintillations of extra-solar point radio sources.

6.1. The inner scale of the turbulent wavenumber spectrum

As noted in Section 2, $\overline{q \epsilon^2} R_\odot$ is mostly determined by density fluctuations near the inner scale q_i^{-1} , where q_i is the boundary between the inertial and dissipative ranges of the wavenumber spectrum $S(q)$ (Equation (2)). Observations of the inner turbulence scale thus allow to infer the magnitude of the spectrum-weighted mean wavenumber of density fluctuations $\overline{q \epsilon^2} R_\odot$.




Figure 7. *Left:* Measured turbulence inner scales q_i^{-1} (Yakovlev et al. 1980; Scott et al. 1983; Coles & Harmon 1989; Manoharan 1993b; Anantharamaiah et al. 1994; Yamauchi et al. 1996, 1998), and magnetic field fluctuations (Bruno & Trenchi 2014; Chiba et al. 2022; Lotz et al. 2023), together with theoretical profiles of key plasma parameters: the proton inertial length $d_i = c/\omega_{pi}$, the proton gyroradius $\rho_i = v_{Ti}/\omega_{ci}$, the proton resonance distance $d_r = (v_{Ti} + v_A)/\omega_{ci}$, calculated using the plasma and magnetic field parameters in Appendix A. The dotted curve shows the relation $q_i^{-1} = 2.5 \times 10^4 (r/R_\odot - 1)^{-1.3}$ cm and the dashed line shows the inner scale model $q_i^{-1} = 10^5 r/R_\odot$ cm by Coles & Harmon (1989). *Right:* The same as the left panel, but for the inner scale wavenumber q_i .

Figure 7 summarizes the pertinent observations of this scale, together with the behavior of several plasma parameters as a function of distance from the Sun. These parameters include the proton inertial length $d_i = c/\omega_{pi}$, the proton gyroradius $\rho_i = v_{Ti}/\omega_{ci}$, the the resonance distance $d_r = (v_{Ti} + v_A)/\omega_{ci}$ for parallel-propagating Alfvén waves (Leamon et al. 1998; Bruno & Trenchi 2014), the proton thermal velocity $v_{Ti} = \sqrt{2k_B T_i/m_p}$, the proton plasma frequency $\omega_{pi} = \sqrt{4\pi n e^2/m_p}$, the proton gyrofrequency $\omega_{ci} = eB/m_p c$, and the Alfvén speed $v_A = B/\sqrt{4\pi n m_p}$. The density, temperature, and magnetic field profiles used to construct these parameter variations are given in Appendix A. Over most of the range from the lower corona ($0.1 R_\odot$) to 1 au and beyond, the inner scale q_i^{-1} is more consistent with the Alfvén wave resonance distance rather than either the proton gyroradius or the proton inertial length. Coles & Harmon (1989) suggest that the inner scale between 2 and $60 R_\odot$ is close to $3c/\omega_{ci}$ and that the data can be approximated by the relation $q_i^{-1} = 10^5 r/R_\odot$ (cm) (see Figure 7). Near 1 au ($r \simeq 200 R_\odot$), the gyroradius ρ_i is smaller than the resonant scattering length d_r , so that finite gyroradius effects dominate the scattering process. Observations by Šafránková et al. (2015) suggest that the location of the density break point between these two domains is controlled by the gyrostructure frequency near 1 au.

6.2. Observations of Extra-Solar Radio Point Sources

Reports of radio observations of extra-solar radio point sources (Rickett 1977; Coles & Harmon 1989; Sakurai & Spangler 1994) typically provide values of the quantity $C_n^2(r)$, which is the normalization constant of the Kolmogorov density power spectrum:

$$\Phi(r; q) = C_n^2(r) q^{-11/3} \quad (21)$$




Figure 8. *Left:* Normalization coefficient $C_n^2(r)$ of the density spectrum given by Equation (23), with $q_i = \omega_{ci}/(v_{Ti} + v_A)$ (see Figure 7). As in Figure 6, the solid/dashed lines are for anisotropy factors $\alpha = 0.25$ (fundamental) and 0.42 (harmonic), respectively, and the grey region denotes the range from $[1/2, 2] \times \overline{q \epsilon^2} R_\odot$. The data from [Coles & Harmon \(1989\)](#) are calculated using the values of $\overline{q \epsilon^2} R_\odot$, with q_i inferred from the break at the larger wavenumber of their density spectrum. Their grey points show where the inner scale was not well defined in their data. The data from [Marsch & Tu \(1990\)](#) are found via a power law fit of the form $A q^{-5/3}$ to the density spectra in their Figure 6, where the slope corresponds to a power law index of $-5/3$, and $C_n^2(r) = A n^2(r)/4\pi$, with n taken from Equation (A1). The dotted line shows the relation $C_n^2(r) = 3.5 \times 10^3 (r/R_\odot - 1)^{-4.7}$. *Right:* As in the left panel, but divided by n^2 , with n given by Equation (A1).

between the outer and inner scales $q_o < q < q_i$, so that⁵ $\delta n^2(r) = \int \Phi(r; q) d^3 q$. Noting that $q_o \ll q_i$, one finds by comparing Equations (1), (2), (9), and (21) that the scaling quantity $C_n^2(r)$ ($\text{cm}^{-20/3}$) is related to the $\overline{q \epsilon^2} R_\odot$ profile by

$$C_n^2(r) = \overline{q \epsilon^2} R_\odot \frac{q_i^{-1/3}}{12 \pi R_\odot} n^2(r), \quad (22)$$

where $n(r)$ is the plasma density. Including the dissipative range above q_i increases $\overline{q \epsilon^2} R_\odot$ by a factor of $\sim 5/3$ due to power-law in dissipative range (see Equation (3)), so that a more appropriate expression, applicable to the wavenumber spectrum in Equation (2), is

$$C_n^2(r) = \overline{q \epsilon^2} R_\odot \frac{q_i^{-1/3}}{20 \pi R_\odot} n^2(r). \quad (23)$$

This $C_n^2(r)$ profile is plotted in the left panel of Figure 8, with the right panel showing the quantity $C_n^2/n^2(r) = \overline{q \epsilon^2} R_\odot q_i^{-1/3}/20 \pi R_\odot$. We note that the right side of Equation (23) depends weakly ($\propto q_i^{-1/3}$) on the inner scale wavenumber, so that even if the inner scale is uncertain to within a factor of two, the value of C_n^2 changes by only 26%.

In addition to the model profile (23), Figure 8 shows values of C_n^2 deduced from observations of scintillations of extra-solar sources ([Spangler et al. 2002](#)). The C_n^2 values from [Spangler et al. \(2002\)](#) appear to be significantly smaller than those in our simulations and in the in-situ observations at 0.4 and 0.8 au by [Marsch & Tu \(1990\)](#). Since Type III solar radio bursts are generated by electrons from active regions/flares and thus propagate not far from the ecliptic (e.g., [Musset et al. 2021](#)), our measurements are more consistent with the larger C_n^2 values from the slow dense solar wind (see similar discussion and conclusion by [Spangler et al. 2002](#)). [Sasikumar Raja et al. \(2016\)](#) suggest that solar cycle variations of C_n^2 might contribute to variations between observations at different times.

⁵ note the $(2\pi)^3$ difference between this normalization and that of Equation (1)




Figure 9. *Left:* Amplitude of the inner-scale density fluctuations $\langle \delta n_i^2 \rangle$ calculated from Equation (25). The grey regions are the same as in Figure 6. The dotted line shows the relation $\langle \delta n_i^2 \rangle = 2 \times 10^7 (r/R_\odot - 1)^{-3.7}$. The inner scale for Coles & Harmon (1989) is taken from the start of the dissipation range visible in their model spectra, and the grey points denote where the inner scale was not well observed in their data. The Marsch & Tu (1990) values are calculated using Equation (24); for their results the inner scale is not observed and instead we use the resonant distance d_r as the measure of the inner scale. As in Figure 6, the black and red data points correspond to the assumption of fundamental and harmonic emission, respectively. *Right:* As in the left panel, but divided by n^2 , with $n(r)$ given by Equation (A1).

The profiles of $\overline{q \epsilon^2} R_\odot$ (Equation (10)) and of the inner scale $q_i^{-1}(r)$ (from Figure 7) also allows us to estimate the amplitude of the density fluctuations at the dominant inner scale. Following, e.g., Chandran et al. (2009) and Sasikumar Raja et al. (2016), we define the squared fractional density perturbation amplitude at the inner-scale wavenumber q_i by

$$\frac{\langle \delta n_i^2 \rangle}{n^2} = 4\pi q_i^3 \frac{S(q_i)}{(2\pi)^3} = 4\pi q_i^3 \frac{\Phi(q_i)}{n^2(r)} = 4\pi q_i^{-2/3} \frac{C_n^2(r)}{n^2(r)}, \quad (24)$$

which also relates the various wavenumber spectrum normalizations $\langle \delta n_i^2 \rangle / n^2$ (Equation (1)), $S(q_i)$ (Equation (2)), and C_n^2 (Equation (21)). One also finds by using Equation (3) that the inner-scale fractional density fluctuation is given by

$$\frac{\langle \delta n_i^2 \rangle}{n^2(r)} = \frac{\overline{q \epsilon^2} R_\odot}{\left(3 + \frac{1}{\delta - 2}\right) q_i(r) R_\odot} \simeq \frac{\overline{q \epsilon^2} R_\odot}{5 q_i(r) R_\odot}, \quad (25)$$

which can be evaluated using the turbulence profile of Equation (10) and the form of $q_i^{-1}(r)$ from Figure 7. This, plus the density profile from Equation (A1), gives the profile of the inner-scale squared density fluctuation values $\langle \delta n_i^2 \rangle$ (cm^{-6}). This quantity, and its dimensionless fractional value $\langle \delta n_i^2 \rangle / n^2$, are shown as functions of $(r/R_\odot - 1)$ in Figure 9.

7. DISCUSSION AND SUMMARY

We have constructed a density fluctuation model for the solar corona and heliosphere that allows quantitative analysis of radio-wave propagation in a medium that is characterized by an anisotropic density turbulence, symmetric around the direction of magnetic field $\vec{\mathbf{B}}$. The density turbulence is characterized by two parameters: the spectrum-weighted mean wavenumber of density fluctuations $\overline{q \epsilon^2}$ (Equation (9)) and the anisotropy measure $\alpha = q_{\parallel}/q_{\perp}$ (Equation (4)). We allow $\overline{q \epsilon^2}$ to be a function of solar distance r from the low corona ($\sim 1.1 R_\odot$) to $215 R_\odot$ (1 au), but the anisotropy parameter α is, for simplicity, considered a constant. The inferred profile of the dimensionless quantity $\overline{q \epsilon^2} R_\odot$ is found to have a broad maximum with a value of about 100 located at around $\sim (4-7) R_\odot$, where the slow solar wind becomes supersonic (Sittler & Guhathakurta 1999). Intriguingly, this is also the range of solar distances where Type III bursts

are observed to have the highest radio spectral flux density (note the broad peak near 1–2 MHz; Sasikumar Raja et al. 2022). The density turbulence model allows quantitative analysis of radio-wave scattering that in turn could be used to decouple the intrinsic properties of solar radio bursts and the effects of radio-wave propagation.

Analysis of the radial variation of the inner scale (i.e., the length associated with the smallest eddies in the inertial range of the turbulence spectrum) with solar distance presents a rather coherent picture. Radio measurements of density inner scales (e.g., Coles & Harmon 1989) are found to be in good agreement with the inner scales deduced from magnetic fluctuations (e.g., Lotz et al. 2023), supporting a close relation between magnetic fluctuations and density fluctuations. It has been argued that kinetic Alfvén waves are a compressive phenomenon that is responsible for both density, magnetic field, and parallel electric field fluctuations near the break between inertial and dissipation ranges (e.g., Chandran et al. 2009; Bian et al. 2010). Over a wide range of distances, from the low solar corona into the solar wind, the turbulence inner scale is comparable (within a factor of two) with the scale of the resonant condition for protons $(v_{Ti} + v_A)/\omega_{ci}$, which is similar to the scale of the break in the spectra of magnetic fluctuations (Bruno & Trenchi 2014; Lotz et al. 2023). Analysis of the inner scales (Coles & Harmon 1989) at (10–50) R_\odot suggests that the proton inertial length $c/\omega_{pi} = v_A/\omega_{ci}$ is a good approximation, while v_{Ti}/ω_{pi} correlates with the break near 1 au (Šafránková et al. 2015). Since $(v_{Ti} + v_A)/\omega_{ci}$ approaches v_{Ti}/ω_{pi} at $r > 50 R_\odot$ and $(v_{Ti} + v_A)/\omega_{ci}$ is dominated by v_A/ω_{ci} at $r > 50 R_\odot$, the resonance condition for protons captures both sets of observations.

Figure 6 shows the profiles $\overline{q\epsilon^2} R_\odot$ inferred from solar, non-solar and in-situ density fluctuation measurements; an acceptable fit to the observations requires the anisotropy parameter to have a value in the range $\alpha \simeq 0.2 - 0.4$. Unfortunately, observations of Type III burst sizes and decay times do not systematically identify whether the observed emission is at the fundamental frequency or its (second) harmonic, and while this ambiguity does affect the best-fit value of the anisotropy parameter α , it has a much smaller effect on the inferred values of $\overline{q\epsilon^2} R_\odot$. Assuming that the sources correspond to emission at the fundamental suggests $\alpha \simeq 0.25$; however, if the emission is at the harmonic then the anisotropy parameter is closer to 0.4. Assuming harmonic emission also requires a factor-of-two larger value of $\overline{q\epsilon^2} R_\odot$, due to the α -dependence in Equation (10); see solid and dashed lines in Figure 6. Since the scattering rate parallel to the magnetic field is a factor $\alpha^2 \ll 1$ times the perpendicular scattering rate (Equation (7)), the radio waves escape predominantly parallel to the magnetic field, so that the degree of anisotropy chiefly affects the duration and the decay time of the bursts, while the source size is mostly controlled by the amplitude of the density fluctuations. The high accuracy of the decay times of Type III bursts allows us to place rather significant constraints on the anisotropy parameter; for example, any value $\alpha > 0.5$ is inconsistent with the available data.

Comparing the results of numerical simulations with observations of source sizes and time profiles, over the range of frequencies $\sim 0.1 - 300$ MHz, we find that the turbulence profile $\overline{q\epsilon^2} R_\odot$ is, within a factor of two, well represented by the empirical form Equation (10). The spatial variation of the fractional density fluctuation amplitude at the inner scale increases monotonically with heliocentric distance, with a noticeable “knee” near the location corresponding to the maximum in $\overline{q\epsilon^2} R_\odot$. The amplitude of the Kolmogorov (1941) density turbulence spectrum varies with distance r from the Sun as $C_n^2(r) \simeq 3.5 \times 10^3 (r/R_\odot - 1)^{-4.7} \text{ cm}^{-20/3}$. The quantity C_n^2 matches well with in-situ density fluctuation measurements at 0.4 and 0.8 au for the slow solar wind and is generally above the measurements of the fast solar wind (Marsch & Tu 1990). However, only the largest values of C_n^2 inferred from the slow solar wind scintillation measurements of Spangler et al. (2002) are in agreement with our model, possibly because Type III bursts propagate mostly in the ecliptic and the scintillation measurements are mostly taken away from the ecliptic. The normalized quantity C_n^2/n^2 demonstrates two distinct domains: it is a constant (or close to a constant) for distances $> 10 R_\odot$, supporting the conclusions of Spangler (2002), but exhibits a growth of three orders of magnitude close to the Sun, reaching a maximum value near (5–7) R_\odot .

The variance of density fluctuations decreases with solar distance approximately as $\langle \delta n_i^2 \rangle(r) = 2 \times 10^7 (r/R_\odot - 1)^{-3.7} \text{ cm}^{-6}$. The exponent in this expression is close to 4 and therefore is not inconsistent with the $\langle \delta n_i^2 \rangle \propto r^{-4}$ expectation (Zank et al. 2017; Tasnim et al. 2022) associated with nearly-incompressible magnetohydrodynamic turbulence. Comparison of the model with observations does, however, suggest that a photosphere-rooted scaling (with a dependence on the quantity $(r/R_\odot - 1)$) could be a better rough symmetry than a simple heliocentric symmetry (i.e., $\propto r^{-4}$), probably due to the turbulence evolution along flux tubes rooted into the photosphere.

The density fluctuations model presented in Equation (10) provides a unifying picture for interpretation of solar radio bursts, extra-solar source broadening, and in-situ measurements of density fluctuations in the solar wind. As the model extends to 1 au, we can predict (Appendix C.2; Equation (C19)) the frequency spectrum and compare with those measured by spacecraft in the slow solar wind. For a 1 au value $\theta_B \simeq 45^\circ$, the ratio of the spectral break




Figure 10. Frequency spectrum of density fluctuations (fundamental is solid, harmonic is dashed line) in accordance with the density fluctuation model, using Equation (C19) with the values of $\langle \delta n_i^2 \rangle / n^2$ at 1 au taken from Figure 9. The grey area shows the [1/2,2] multiplication-factor interval used in the previous figures. The in-situ fluctuations data at 1 au (Celnikier et al. 1983, 1987; Kellogg & Horbury 2005; Chen et al. 2012, 2014) are shown for comparison.

frequency \tilde{f}_i to its value f_i for an isotropic ($\alpha = 1$) spectrum is $\sqrt{0.5(1 + \alpha^2)}$ (see remark after Equation (C19)), which is only weakly sensitive to α in the range 0.25 – 0.4, so that the spectral break point is about the same for all α values in this range. Figure 10 demonstrates good agreement between the model and observations near the break of the spectrum at $f \simeq f_i$. The discrepancy below the break frequency is likely to be associated with flattening of the density fluctuation spectra due to kinetic Alfvén waves (Chandran et al. 2009) and has been observed close to the Sun before (e.g., Coles & Harmon 1989). As discussed, this flattening is not included into our model, but could be used for further improvement of the model.

An important feature of the model is that it provides a quantitative description for solar burst source sizes, apparent positions, and decay times, with the results summarized below:

Source sizes: Type III burst source sizes are predominantly determined by radio-wave scattering over the entire range of frequencies and follow a $1/f$ trend (both in the simulations and observations) from 300 MHz down to 0.1 MHz. The source sizes do not depend on burst intensity (e.g. Saint-Hilaire et al. 2013) and the source sizes for fine structures in Type III bursts and Type III spikes are rather similar (e.g., Kontar et al. 2017; Clarkson et al. 2023). At frequencies below 0.1 MHz, the source sizes exceed 120 degrees (see Figure 2), comparable to half the sky, so that the source effectively surrounds the observer making unambiguous measurements more challenging.

Source positions: From Figure 5 we see that for high frequency sources between 100 and 300 MHz the apparent source positions for both fundamental ($\omega = \omega_{pe}$) and harmonic ($\omega = 2\omega_{pe}$) emission are at a location corresponding to emission at double the plasma frequency. However, for emission frequencies from ~ 100 MHz down to ~ 0.1 MHz, the apparent source location is further away from the Sun than the location corresponding to emission at the plasma frequency or even to the location corresponding to emission at double the plasma frequency. This result elegantly resolves the long-standing conundrum that source positions observed at the fundamental and the harmonic are virtually co-spatial (Suzuki & Dulk 1985): the apparent source position is moved away from the Sun by scattering effects, which are largely independent of whether fundamental or harmonic emission is involved. At low frequencies $\lesssim 0.2$ MHz, this effect is even more pronounced: the source positions are shifted by 0.5 au, so that an observer at 1 au is actually embedded in the apparent source.

Type III burst decay times: The decay time of Type III solar radio bursts is well approximated by an inverse-frequency ($1/f$) dependence between 0.1 and 300 MHz. Our scattering simulations reveal a rather intricate picture, in which scattering serves to provide a fundamental “floor” on the observed duration of radio bursts emitted via plasma emission. Below 1 MHz, the average Type III burst decay time is consistent with the scattering simulations, but at frequencies above a few MHz, the typical Type III burst decay time is longer than the simulated value, suggesting that the decay time for such bursts is determined by processes that operate on timescales longer than scattering, such as electron propagation, electron injection, and/or intrinsic emission time scales. The shortest features in the dynamic spectrum, such as Type IIIb striae (e.g., Kontar et al. 2017; Sharykin et al. 2018), drift pairs (Kuznetsov & Kontar 2019), spikes (Clarkson et al. 2021), and fine structures in type II solar radio bursts (Chrysaphi et al. 2018) are all consistent with the scattering model. Statistical analysis of ~ 1100 fine structures (Clarkson et al. 2023) near 30 MHz, with 0.02 s resolution, indeed shows that there are virtually no bursts shorter than 0.2 s and that the typical burst decay time is about 0.3 s, consistent with the simulations in Figure 3.

In summary, solar burst shortest time profiles, source sizes, and positions are determined mainly by propagation effects (mostly anisotropic scattering) and not by intrinsic properties of the radio source. A detailed knowledge of the scattering process paves the way to disentangling scattering effects from observations and so better constraining the intrinsic properties of solar radio burst sources. Since individual source sizes and decay times are typically measured more accurately than the spread in measurements of multiple sources, it is therefore likely that the spread in observational properties is due to varying levels of turbulence and plasma density in different events. Varying the magnitude of the $\overline{q\epsilon^2} R_\odot$ profile (Equation (10)) by a factor in the range 0.5 to 2 covers the majority of the observations, while extending this multiplicative factor by a further factor of two (to a range between 0.25 to 4) covers virtually all observed data points (except for some extreme outliers). We thus conclude that the $\overline{q\epsilon^2} R_\odot$ profile of Equation (10) is variable within a factor of about two. The broadening of extra-solar point sources by the turbulent solar atmosphere and solar radio burst measurements are complementary data sets. We note the considerable data gap between ground-based and space-based solar burst observations in the range 3–20 MHz (where extra-solar observations appear essential), and encourage the development of observations to fill this gap and hence further constrain the level of turbulence in the inner heliosphere. The Sun Radio Interferometer Space Experiment (SunRISE) is likely to provide much needed data.

ACKNOWLEDGMENTS

We thank P. Massa for helpful discussions. EPK, DLC and XC acknowledge financial support from the STFC Consolidated Grant ST/T000422/1. AGE was supported by NASA Kentucky under NASA award number 80NSSC21M0362. NC acknowledges funding support from the Initiative Physique des Infinis (IPI), a research training program of the IDEX SUPER at Sorbonne Université. We also acknowledge support from the International Space Science Institute for the LOFAR <http://www.issibern.ch/teams/lofar/> and solar flare <http://www.issibern.ch/teams/solflareconnectsolenerg/teams>.

APPENDIX

A. PLASMA DENSITY, MAGNETIC FIELD AND TEMPERATURE PROFILES

The density profile $n(r)$ (cm^{-3}) used for the solar burst scattering simulations by Kontar et al. (2019), Kuznetsov et al. (2020), and Chen et al. (2020, 2023) is

$$n(r) = 4.8 \times 10^9 \left(\frac{R_\odot}{r} \right)^{14} + 3 \times 10^8 \left(\frac{R_\odot}{r} \right)^6 + 1.4 \times 10^6 \left(\frac{R_\odot}{r} \right)^{2.3}. \quad (\text{A1})$$

This density profile (i) closely follows the Parker (1958) density model, (ii) has an analytical form that can be easily differentiated to find the derivatives that are useful in solving the ray-tracing equations, (iii) is consistent at low heights with the higher densities observed in flaring active regions that produce Type III bursts (e.g., Holman et al.

2011; Reid et al. 2014), and (iv) is suitable for Type III burst modelling in the solar corona and heliosphere (Mann et al. 1999; Kontar 2001; Reid & Kontar 2021). The left panel of Figure 11 shows the model profile compared with different density models frequently used in the literature.




Figure 11. *Left panel:* Coronal density model (black solid line) given by Equation (A1). Other coronal density models (Allen 1947; Newkirk 1961; Alvarez & Haddock 1973b; Saito et al. 1977; Bougeret et al. 1984b; Bird et al. 1994; Leblanc et al. 1998; Mancuso & Spangler 2000) are also shown, within the ranges of their respective validity. The models from Bird et al. (1994) and Mancuso & Spangler (2000) are as used in Spangler (2002). *Right panel:* Magnetic field model (black solid line) given by Equation (A2), together with observations (Burlaga & Ness 1968; Villante & Mariani 1975; Behannon 1976; Rosenberg et al. 1978; Dulk & McLean 1978; Patzold et al. 1987; Sakurai & Spangler 1994; Lin et al. 2000; Spangler 2005; Ramesh et al. 2010; Gopalswamy et al. 2012; Kolotkov et al. 2018; Spangler 2020). The red line shows the model of Dulk & McLean (1978)

The magnetic field $B(r)$ (G) model, shown in the right panel of Figure 11, is given by

$$B(r) = 0.5 \frac{\left(\frac{r}{R_{\odot}} - 1\right)^{-1.5}}{\left(\frac{r}{10 R_{\odot}} + 1\right)} + 1.18 \left(\frac{R_{\odot}}{r}\right)^2, \quad (\text{A2})$$

where the first term is introduced to accommodate the results of Dulk & McLean (1978), applicable for near-solar distances $r \lesssim 10 R_{\odot}$, into the form of the interplanetary magnetic field for $r \gg 10 R_{\odot}$ (see discussion by Patzold et al. 1987).

The ion temperature T_i (K) is given by

$$T_i(r) = 2.4 \times 10^6 \left(\frac{r}{10 R_{\odot}} + 1\right)^{-0.74} \text{ K}, \quad (\text{A3})$$

where the interplanetary dependency $r^{-0.74}$ (with $T_i = 2.5 \times 10^5$ K at 1 au = $215 R_{\odot}$) is taken from Hellinger et al. (2011) and has been adjusted to accommodate the range $r \lesssim 10 R_{\odot}$ in accordance with the results of Dulk & McLean (1978).

B. ANGULAR BROADENING OF EXTRA-SOLAR POINT SOURCES

The overall angular broadening of a point source at infinity can be calculated by integrating the angular broadening rate along the path (z -direction) traversed by the radio waves. For the most significant broadening, i.e., that along

the \perp_2 direction perpendicular to the (radial) magnetic field in the plane of the sky, the rate of diffusive broadening is given by Equation (7), repeated here:

$$\frac{d\langle\theta_{\perp}^2\rangle}{dz} = \frac{\pi}{8} \frac{\omega_{pe}^4}{\omega^4} \frac{1}{q\epsilon^2} .$$

If r is the distance of closest approach to the Sun of a ray from a distance point source (Figure 2), then the apparent source, observed at 1 au, will be broadened by an amount

$$\langle\theta_{\perp}^2\rangle = \frac{\pi}{8\omega^4} \int_{-\infty}^{1 \text{ au}} \omega_{pe}^4 \left(\sqrt{r^2+z^2}\right) \frac{1}{q\epsilon^2} \left(\sqrt{r^2+z^2}\right) dz = \frac{2\pi^3 e^4}{m_e^2 \omega^4} \int_{-\infty}^{1 \text{ au}} n^2 \left(\sqrt{r^2+z^2}\right) \frac{1}{q\epsilon^2} \left(\sqrt{r^2+z^2}\right) dz , \quad (\text{B4})$$

where we have used $\omega_{pe} = \sqrt{4\pi n e^2/m_e}$. With the substitution $z = r \tan \chi$, this can be written

$$\langle\theta_{\perp}^2\rangle = \alpha \frac{2\pi^3 e^4}{m_e^2 \omega^4} \left(\frac{r}{R_{\odot}}\right) \int_{-\pi/2}^{\tan^{-1}(215 R_{\odot}/r)} n^2(r \sec \chi) \frac{1}{q\epsilon^2} R_{\odot}(r \sec \chi) \sec^2 \chi d\chi . \quad (\text{B5})$$

Equation (B5) provides the formal expression corresponding to the approximate proportionalities in Equation (8). Using the form of $\frac{1}{q\epsilon^2} R_{\odot}$ from our nominal model (10) and the density model from Equation (A1), we obtain

$$\begin{aligned} \langle\theta_{\perp}^2\rangle = & 2 \times 10^3 \alpha \times \left(\frac{e^4}{8\pi m_e^2 f^4}\right) \left(\frac{r}{R_{\odot}}\right)^{0.3} \times \\ & \int_{-\pi/2}^{\tan^{-1}(215 R_{\odot}/r)} \left[4.8 \times 10^9 \left(\frac{R_{\odot}}{r}\right)^{14} \cos^{14} \chi + 3 \times 10^8 \left(\frac{R_{\odot}}{r}\right)^6 \cos^6 \chi + 1.4 \times 10^6 \left(\frac{R_{\odot}}{r}\right)^{2.3} \cos^{2.3} \chi \right]^2 \times \\ & \times \left(1 - \frac{R_{\odot} \cos \chi}{r}\right)^{2.7} \frac{d\chi}{\cos^{1.3} \chi} . \end{aligned} \quad (\text{B6})$$

At large closest-approach distances $r \gg R_{\odot}$, the density profile (A1) is well approximated by the last term in the square brackets, i.e., by the simple power-law $n(r) \propto r^{-2.3}$, and the term $(1 - R_{\odot} \cos \chi/r)^{2.7} \simeq 1$. In this situation, the FWHM size assumes the relatively simple form

$$\text{FWHM}_{\perp} = 2.35 \sqrt{\alpha} \times \sqrt{2 \times 10^3} \times (1.4 \times 10^6) \times \left[\frac{\text{B}(1; 2.15, 0.5) + \text{B}(\eta; 2.15, 0.5)}{16\pi} \right]^{1/2} \left(\frac{e^2}{m_e}\right) \left(\frac{r}{R_{\odot}}\right)^{-2.15} \times \frac{1}{f^2} , \quad (\text{B7})$$

where $\text{B}(\eta; u, v)$ are the (incomplete if $\eta < 1$) beta functions corresponding to the integrals $2 \int_{-\pi/2}^0 \cos^{3.3} \chi d\chi$ and $2 \int_0^{\tan^{-1}(215 R_{\odot}/r)} \cos^{3.3} \chi d\chi$, respectively, and $\eta = [1 + (r/215 R_{\odot})^2]^{-1}$. Scaling to the nominal frequency $f = 1.5$ GHz (Figure 2), this evaluates to

$$\text{FWHM}_{\perp} \simeq 4 \times 10^{-3} \sqrt{\alpha} \left(\frac{r}{R_{\odot}}\right)^{-2.15} \left(\frac{1.5 \text{ GHz}}{f}\right)^2 \text{ radians} \simeq 800 \sqrt{\alpha} \left(\frac{r}{R_{\odot}}\right)^{-2.15} \left(\frac{1.5 \text{ GHz}}{f}\right)^2 \text{ arcseconds} . \quad (\text{B8})$$

Equation (B8) provides a simple, but accurate, analytical approximation for FWHM_{\perp} , valid for $r \gg R_{\odot}$.

C. FORMS OF THE IN SITU DENSITY FLUCTUATION FREQUENCY SPECTRA

The in situ frequency power spectrum $P(f)$ of density fluctuations measured by a spacecraft is related to the wavenumber spectrum $S(\vec{q})$ at a single location in the solar wind frame through the relation (11), repeated here:

$$P(f) = \frac{n^2}{(2\pi)^3} \int S(\vec{q}) \delta\left(\frac{\vec{q} \cdot \vec{V}_{\text{SW}}}{2\pi} - f\right) d^3 q .$$

In this Appendix we explore the forms of $P(f)$ for different forms of the wavenumber spectrum $S(\vec{q})$.

C.1. *Isotropic Wavenumber Spectrum*

Our first example is the simplest case of an isotropic spectrum: $S(\vec{q}) = S(q)$. For such a case, Equation (11) may be written

$$\begin{aligned} P(f) &= \frac{n^2}{(2\pi)^2} \int_{q_{\perp}=0}^{\infty} \int_{q_{\parallel}=-\infty}^{\infty} S\left(\sqrt{q_{\parallel}^2 + q_{\perp}^2}\right) \delta\left(\frac{q_{\parallel} V_{\text{SW}}}{2\pi} - f\right) q_{\perp} dq_{\perp} dq_{\parallel} \\ &= \frac{n^2}{2\pi V_{\text{SW}}} \int_{q_{\perp}=0}^{\infty} S\left(\sqrt{q_{\perp}^2 + \left(\frac{2\pi f}{V_{\text{SW}}}\right)^2}\right) q_{\perp} dq_{\perp} , \end{aligned} \quad (\text{C9})$$

Here, we evaluate this for a specific form of $S(q)$, viz. the broken power-law around q_i (cf. Equation (2)), with an outer scale $q_o^{-1} \gg q_i^{-1}$, so that the contribution to $q \epsilon^2 R_{\odot}$ from the large scales $> q_o^{-1}$ is negligible:

$$S(q) = S(q_i) \times \begin{cases} \left(\frac{q}{q_i}\right)^{-11/3} ; & q \leq q_i \\ \left(\frac{q}{q_i}\right)^{-2-\delta} ; & q > q_i . \end{cases} \quad (\text{C10})$$

Evaluation of the fluctuation frequency spectrum $P(f)$ requires consideration of two domains:

$$(i) f \leq f_i = \frac{q_i V_{\text{SW}}}{2\pi}$$

Here the resonant $q_{\parallel} = 2\pi f/V_{\text{SW}} < q_i$, so that for values of $q_{\perp}^2 < q_i^2 - q_{\parallel}^2$ the lower (inertial) branch of the wavenumber spectrum applies, while for values of $q_{\perp}^2 > q_i^2 - q_{\parallel}^2$ the upper (dissipative) branch of the wavenumber spectrum is relevant. The integral is thus composed of two parts:

$$\begin{aligned} P(f) &= \frac{n^2}{2\pi V_{\text{SW}}} S(q_i) \left[q_i^{11/3} \int_{q_{\perp}=0}^{\sqrt{q_i^2 - (2\pi f/V_{\text{SW}})^2}} \left(q_{\perp}^2 + \left(\frac{2\pi f}{V_{\text{SW}}}\right)^2 \right)^{-11/6} q_{\perp} dq_{\perp} + \right. \\ &\quad \left. + q_i^{\delta+2} \int_{q_{\perp}=\sqrt{q_i^2 - (2\pi f/V_{\text{SW}})^2}}^{\infty} \left(q_{\perp}^2 + \left(\frac{2\pi f}{V_{\text{SW}}}\right)^2 \right)^{-1-\frac{\delta}{2}} q_{\perp} dq_{\perp} \right] \\ &= \frac{n^2}{4\pi^2 f_i} q_i^3 S(q_i) \left[\frac{3}{5} \left(\left(\frac{2\pi f}{q_i V_{\text{SW}}}\right)^{-5/3} - 1 \right) + \frac{1}{\delta} \right] \\ &= \frac{\langle \delta n_i^2 \rangle}{2 f_i} \left[\frac{3}{5} \left(\left(\frac{f}{f_i}\right)^{-5/3} - 1 \right) + \frac{1}{\delta} \right] , \end{aligned} \quad (\text{C11})$$

where in the last equality we have used Equation (24).

$$(ii) f > f_i = \frac{q_i V_{\text{SW}}}{2\pi}$$

Here the resonant $q_{\parallel} = 2\pi f/V_{\text{SW}} > q_i$, so that for all values of q_{\perp} only the upper branch of the wavenumber spectrum is relevant. Thus

$$\begin{aligned} P(f) &= \frac{n^2}{2\pi V_{\text{SW}}} S(q_i) q_i^{\delta+2} \int_{q_{\perp}=0}^{\infty} \left(q_{\perp}^2 + \left(\frac{2\pi f}{V_{\text{SW}}}\right)^2 \right)^{-1-\frac{\delta}{2}} q_{\perp} dq_{\perp} \\ &= \frac{n^2}{4\pi^2 f_i} q_i^3 S(q_i) \frac{1}{\delta} \left(\frac{2\pi f}{q_i V_{\text{SW}}}\right)^{-\delta} \\ &= \frac{\langle \delta n_i^2 \rangle}{2 f_i} \frac{1}{\delta} \left(\frac{f}{f_i}\right)^{-\delta} \end{aligned} \quad (\text{C12})$$

Combining Equations (C11) and (C12),

$$P(f) = \frac{\langle \delta n_i^2 \rangle}{2 f_i} \times \begin{cases} \frac{3}{5} \left[\left(\frac{f}{f_i} \right)^{-5/3} - \left(1 - \frac{5}{3\delta} \right) \right] & ; \quad f \leq f_i \\ \frac{1}{\delta} \left(\frac{f}{f_i} \right)^{-\delta} & ; \quad f > f_i . \end{cases} \quad (\text{C13})$$

As a check on this result, we can evaluate (using algebra) the quantity

$$\int_0^\infty f P(f) df = \frac{\langle \delta n_i^2 \rangle}{2 f_i} \left(3 + \frac{1}{\delta - 2} \right) \frac{f_i^2}{2} = \frac{1}{4} \left(3 + \frac{1}{\delta - 2} \right) \langle \delta n_i^2 \rangle f_i . \quad (\text{C14})$$

so that, using the general result (18), with $\alpha = 1$ for an isotropic spectrum, the value of $\overline{q \epsilon^2} R_\odot$ is

$$\overline{q \epsilon^2} R_\odot = \frac{8\pi}{n^2 V_{\text{SW}}} R_\odot \int_0^\infty f P(f) df = \left(3 + \frac{1}{\delta - 2} \right) (q_i R_\odot) \frac{\langle \delta n_i^2 \rangle}{n^2} , \quad (\text{C15})$$

in agreement with Equation (25).

The local power-law index of the frequency power spectrum is the negative of its logarithmic derivative, viz.

$$\delta_{\text{loc}} = - \frac{d \ln P(f)}{d \ln f} \equiv - \frac{f}{P} \frac{dP}{df} = \begin{cases} \frac{5/3}{1 - (1 - \frac{5}{3\delta}) (\frac{f}{f_i})^{5/3}} & ; \quad f \leq f_i \\ \delta & ; \quad f > f_i . \end{cases} \quad (\text{C16})$$

At low frequencies $f \ll f_i$, the frequency spectrum has a power-law index $\delta_{\text{loc}} = 5/3$; as f increases, the spectrum gradually steepens to $\delta_{\text{loc}} = \delta$ as f approaches f_i , and continues with that value thereafter. Thus a double-power law spectrum in wavenumber does not correspond to a double power-law spectrum in frequency, neither does a double-power law in frequency correspond to a double power-law in wavenumber space. Interestingly, the variation in the spectrum of magnetic fluctuations reported by Sioulas et al. (2023, their Fig. 2) near the break frequency f_i , is broadly consistent with Equation (C16).

C.2. Anisotropic density turbulence Spectrum with anisotropic spectral break points

For anisotropic turbulence with a spectral break point that is also anisotropic (see left panel of Figure 12),

$$S(q) = S(q_i) \times \begin{cases} \left(\frac{\sqrt{q_\perp^2 + \frac{q_\parallel^2}{\alpha^2}}}{q_i} \right)^{-11/3} & ; \quad \sqrt{q_\perp^2 + \frac{q_\parallel^2}{\alpha^2}} \leq q_i \\ \left(\frac{\sqrt{q_\perp^2 + \frac{q_\parallel^2}{\alpha^2}}}{q_i} \right)^{-2-\delta} & ; \quad \sqrt{q_\perp^2 + \frac{q_\parallel^2}{\alpha^2}} > q_i . \end{cases} \quad (\text{C17})$$

The basic result (11) can be transformed from the wavevector \vec{q} to the variable $\widetilde{\vec{q}}$ (cf. Equation (12)), giving

$$P(f) = \alpha \frac{n^2}{(2\pi)^3} \int S(\widetilde{\vec{q}}) \delta \left(\frac{\widetilde{\vec{q}} \cdot \widetilde{\vec{V}}_{\text{SW}}}{2\pi} - f \right) d^3 \widetilde{q} , \quad (\text{C18})$$

where $\widetilde{\vec{V}}_{\text{SW}} = (V_\perp, \alpha V_\parallel)$. Since $S(\widetilde{\vec{q}})$ is isotropic, a similar analysis to that of Appendix C.1 yields the spectrum

$$P(f) = \alpha \frac{\langle \delta n_i^2 \rangle}{2 \widetilde{f}_i} \times \begin{cases} \frac{3}{5} \left[\left(\frac{f}{\widetilde{f}_i} \right)^{-5/3} - \left(1 - \frac{5}{3\delta} \right) \right] & ; \quad f \leq \widetilde{f}_i \\ \frac{1}{\delta} \left(\frac{f}{\widetilde{f}_i} \right)^{-\delta} & ; \quad f > \widetilde{f}_i , \end{cases} \quad (\text{C19})$$

where $\widetilde{f}_i = q_i \widetilde{V}_{\text{SW}} / 2\pi$, with $\widetilde{V}_{\text{SW}} = V_{\text{SW}} (\sin^2 \theta_B + \alpha^2 \cos^2 \theta_B)^{1/2}$, θ_B being the angle between the solar wind velocity \vec{V}_{SW} and the magnetic field \vec{B} . The form of the local power-law spectral index δ_{loc} is similar to Equation (C16), but

with f_i replaced by \tilde{f}_i . In the case where the solar wind velocity \vec{V}_{SW} is parallel to the magnetic field \vec{B} , $\tilde{V}_{\text{SW}} = \alpha V_{\text{SW}}$, $\tilde{f}_i = \alpha f_i$, and so

$$P(f) = \frac{\langle \delta n_i^2 \rangle}{2 f_i} \times \begin{cases} \frac{3}{5} \left[\left(\frac{f}{\alpha f_i} \right)^{-5/3} - \left(1 - \frac{5}{3\delta} \right) \right] & ; \quad f \leq \alpha f_i \\ \frac{1}{\delta} \left(\frac{f}{\alpha f_i} \right)^{-\delta} & ; \quad f > \alpha f_i \end{cases} \quad (\text{C20})$$




Figure 12. Forms of the parallel and perpendicular wavenumber spectra corresponding to the two cases described in Appendices C.2 (Equation (C17); left) and C.3 (Equation (C21); right). In the former case, the spectral break points are different for parallel and perpendicular directions, while in the latter case the spectral break points for both directions are the same. In both plots, both axes are on logarithmic scales.

C.3. Anisotropic spectrum with isotropic break point

It has been argued by [Armstrong et al. \(1990\)](#) that a wavenumber spectrum that breaks at a prescribed value of $q = \sqrt{q_{\parallel}^2 + q_{\perp}^2}$ in all directions (despite the anisotropy in the wavenumber distribution) is a better fit to the observations. This corresponds to the following density turbulence spectrum (see right panel of Figure 12):

$$S(\vec{q}) = S(q_i) \times \begin{cases} \left(\frac{\sqrt{q_{\perp}^2 + \frac{q_{\parallel}^2}{\alpha^2}}}{q_i} \right)^{-11/3} & ; \quad \sqrt{q_{\perp}^2 + q_{\parallel}^2} \leq q_i \\ \left(\frac{\sqrt{q_{\perp}^2 + \frac{q_{\parallel}^2}{\alpha^2}}}{q_i} \right)^{-2-\delta} & ; \quad \sqrt{q_{\perp}^2 + q_{\parallel}^2} > q_i \end{cases} \quad (\text{C21})$$

Note that the argument of S reflects the anisotropy in $S(\vec{q})$ and so is different than the isotropic case of Appendix C.1 (Equation (C10)). However, the spectral break point is the same as in Appendix C.1, and so differs from Appendix C.2, where the spectral break point occurs at different values of q for different directions (Equation (C17)).

As before, the resonance occurs when $q_{\parallel} = 2\pi f/V_{\text{SW}}$, and so we have, similar to Equation (C9), but with an additional factor of α appearing in the argument of S ,

$$P(f) = \frac{n^2}{2\pi V_{\text{SW}}} \int_{q_{\perp}=0}^{\infty} S \left(\sqrt{q_{\perp}^2 + \left(\frac{2\pi f}{\alpha V_{\text{SW}}} \right)^2} \right) q_{\perp} dq_{\perp} \quad (\text{C22})$$

The derivation of $P(f)$ proceeds similarly to Appendix C.1, but recognizes the appearance of the factor α in the second term in the argument of S :

(i) $f \leq f_i = \frac{q_i V_{\text{SW}}}{2\pi}$

$$\begin{aligned}
P(f) &= \frac{n^2}{2\pi V_{\text{SW}}} S(q_i) \left[q_i^{11/3} \int_{q_{\perp}=0}^{\sqrt{q_i^2 - (2\pi f/V_{\text{SW}})^2}} \left(q_{\perp}^2 + \left(\frac{2\pi f}{\alpha V_{\text{SW}}} \right)^2 \right)^{-11/6} q_{\perp} dq_{\perp} + \right. \\
&\quad \left. + q_i^{\delta+2} \int_{q_{\perp}=\sqrt{q_i^2 - (2\pi f/V_{\text{SW}})^2}}^{\infty} \left(q_{\perp}^2 + \left(\frac{2\pi f}{\alpha V_{\text{SW}}} \right)^2 \right)^{-1-\frac{\delta}{2}} q_{\perp} dq_{\perp} \right] \\
&= \frac{n^2}{4\pi^2 f_i} q_i^3 S(q_i) \left[\frac{3}{5} \left\{ \left(\frac{2\pi f}{\alpha q_i V_{\text{SW}}} \right)^{-5/3} - \left(1 - \left(\frac{2\pi f}{q_i V_{\text{SW}}} \right)^2 \left(1 - \frac{1}{\alpha^2} \right) \right)^{-5/6} \right\} + \right. \\
&\quad \left. + \frac{1}{\delta} \left(1 - \left(\frac{2\pi f}{q_i V_{\text{SW}}} \right)^2 \left(1 - \frac{1}{\alpha^2} \right) \right)^{-\delta/2} \right] \\
&= \frac{\langle \delta n_i^2 \rangle}{2 f_i} \left[\frac{3}{5} \left\{ \left(\frac{f}{\alpha f_i} \right)^{-5/3} - \left(1 - \left(\frac{f}{f_i} \right)^2 \left(1 - \frac{1}{\alpha^2} \right) \right)^{-5/6} \right\} + \frac{1}{\delta} \left(1 - \left(\frac{f}{f_i} \right)^2 \left(1 - \frac{1}{\alpha^2} \right) \right)^{-\delta/2} \right]. \quad (\text{C23})
\end{aligned}$$

(ii) $f > f_i = \frac{q_i V_{\text{SW}}}{2\pi}$

$$\begin{aligned}
P(f) &= \frac{n^2}{2\pi V_{\text{SW}}} S(q_i) q_i^{\delta+2} \int_{q_{\perp}=0}^{\infty} \left(q_{\perp}^2 + \left(\frac{2\pi f}{\alpha V_{\text{SW}}} \right)^2 \right)^{-1-\frac{\delta}{2}} q_{\perp} dq_{\perp} \\
&= \frac{n^2}{4\pi V_{\text{SW}}} S(q_i) \frac{2}{\delta} q_i^{\delta+2} \left(\frac{2\pi f}{\alpha V_{\text{SW}}} \right)^{-\delta} \\
&= \frac{\langle \delta n_i^2 \rangle}{2 f_i} \frac{1}{\delta} \left(\frac{f}{\alpha f_i} \right)^{-\delta} \quad (\text{C24})
\end{aligned}$$

Results (C23) and (C24) match at $f = f_i$ and also reduce to the isotropic result (C13) when $\alpha = 1$. However, the results (12) through (18) relating $q \epsilon^2 R_{\odot}$ and $\int_0^{\infty} f P(f) df$ do not hold in the general case $\alpha \neq 1$, because the wavenumber spectrum $S(\tilde{q})$ in Equation (C21) is not, and cannot be reduced to, an isotropic form.

REFERENCES

- Abranin, E. P., Bazelian, L. L., Goncharov, N. I., et al. 1976, *Soviet Ast.*, 19, 602
- . 1978, *SoPh*, 57, 229, doi: [10.1007/BF00152056](https://doi.org/10.1007/BF00152056)
- Alexander, J. K., Malitson, H. H., & Stone, R. G. 1969, *SoPh*, 8, 388, doi: [10.1007/BF00155385](https://doi.org/10.1007/BF00155385)
- Alexandrova, O., Chen, C. H. K., Sorriso-Valvo, L., Horbury, T. S., & Bale, S. D. 2013, *SSRv*, 178, 101, doi: [10.1007/s11214-013-0004-8](https://doi.org/10.1007/s11214-013-0004-8)
- Allen, C. W. 1947, *MNRAS*, 107, 426, doi: [10.1093/mnras/107.5-6.426](https://doi.org/10.1093/mnras/107.5-6.426)
- Alvarez, H. 1976, *SoPh*, 46, 483, doi: [10.1007/BF00149873](https://doi.org/10.1007/BF00149873)
- Alvarez, H., & Haddock, F. T. 1973a, *SoPh*, 30, 175, doi: [10.1007/BF00156186](https://doi.org/10.1007/BF00156186)
- . 1973b, *SoPh*, 29, 197, doi: [10.1007/BF00153449](https://doi.org/10.1007/BF00153449)
- Anantharamaiah, K. R., Gothoskar, P., & Cornwell, T. J. 1994, *Journal of Astrophysics and Astronomy*, 15, 387, doi: [10.1007/BF02714823](https://doi.org/10.1007/BF02714823)
- Armand, N. A., Efimov, A. I., & Iakovlev, O. I. 1987, *A&A*, 183, 135
- Armstrong, J. W., Coles, W. A., Kojima, M., & Rickett, B. J. 1990, *ApJ*, 358, 685, doi: [10.1086/169022](https://doi.org/10.1086/169022)
- Arzner, K., & Magun, A. 1999, *A&A*, 351, 1165
- Barrow, C. H., & Achong, A. 1975, *SoPh*, 45, 459, doi: [10.1007/BF00158462](https://doi.org/10.1007/BF00158462)
- Barrow, C. H., Zarka, P., & Aubier, M. G. 1994, *A&A*, 286, 597
- Bastian, T. S. 1994, *ApJ*, 426, 774, doi: [10.1086/174114](https://doi.org/10.1086/174114)
- Behannon, K. W. 1976, in *Physics of Solar Planetary Environments*, ed. D. J. Williams, Vol. 1, 332–345
- Bian, N. H., Emslie, A. G., & Kontar, E. P. 2019, *ApJ*, 873, 33, doi: [10.3847/1538-4357/ab0411](https://doi.org/10.3847/1538-4357/ab0411)
- Bian, N. H., Kontar, E. P., & Brown, J. C. 2010, *A&A*, 519, A114+, doi: [10.1051/0004-6361/201014048](https://doi.org/10.1051/0004-6361/201014048)

- Bird, M. K., Volland, H., Paetzold, M., et al. 1994, *ApJ*, 426, 373, doi: [10.1086/174073](https://doi.org/10.1086/174073)
- Blesing, R. G., & Dennison, P. A. 1972, *PASA*, 2, 84, doi: [10.1017/S1323358000012947](https://doi.org/10.1017/S1323358000012947)
- Bougeret, J.-L., Caroubalos, C., Mercier, C., & Pick, M. 1970, *A&A*, 6, 406
- Bougeret, J. L., Fainberg, J., & Stone, R. G. 1984a, *A&A*, 141, 17
- Bougeret, J. L., King, J. H., & Schwenn, R. 1984b, *SoPh*, 90, 401, doi: [10.1007/BF00173965](https://doi.org/10.1007/BF00173965)
- Bougeret, J. L., Goetz, K., Kaiser, M. L., et al. 2008, *SSRv*, 136, 487, doi: [10.1007/s11214-007-9298-8](https://doi.org/10.1007/s11214-007-9298-8)
- Bradford, H. M., & Routledge, D. 1980, *MNRAS*, 190, 73P, doi: [10.1093/mnras/190.1.73P](https://doi.org/10.1093/mnras/190.1.73P)
- Breen, A. R., Mikic, Z., Linker, J. A., et al. 1999, *J. Geophys. Res.*, 104, 9847, doi: [10.1029/1998JA900091](https://doi.org/10.1029/1998JA900091)
- Bruno, R., & Trenchi, L. 2014, *ApJL*, 787, L24, doi: [10.1088/2041-8205/787/2/L24](https://doi.org/10.1088/2041-8205/787/2/L24)
- Burlaga, L. F., & Ness, N. F. 1968, *Canadian Journal of Physics Supplement*, 46, 962, doi: [10.1139/p68-394](https://doi.org/10.1139/p68-394)
- Cecconi, B. 2014, *Comptes Rendus Physique*, 15, 441, doi: [10.1016/j.crhy.2014.02.005](https://doi.org/10.1016/j.crhy.2014.02.005)
- Celnikier, L. M., Harvey, C. C., Jegou, R., Moricet, P., & Kemp, M. 1983, *A&A*, 126, 293
- Celnikier, L. M., Muschietti, L., & Goldman, M. V. 1987, *A&A*, 181, 138
- Chandran, B. D. G., Quataert, E., Howes, G. G., Xia, Q., & Pongkitiwanichakul, P. 2009, *ApJ*, 707, 1668, doi: [10.1088/0004-637X/707/2/1668](https://doi.org/10.1088/0004-637X/707/2/1668)
- Chandrasekhar, S. 1952, *MNRAS*, 112, 475, doi: [10.1093/mnras/112.5.475](https://doi.org/10.1093/mnras/112.5.475)
- Chen, C. H. K., Salem, C. S., Bonnell, J. W., Mozer, F. S., & Bale, S. D. 2012, *PhRvL*, 109, 035001, doi: [10.1103/PhysRevLett.109.035001](https://doi.org/10.1103/PhysRevLett.109.035001)
- Chen, C. H. K., Sorriso-Valvo, L., Šafránková, J., & Němeček, Z. 2014, *ApJL*, 789, L8, doi: [10.1088/2041-8205/789/1/L8](https://doi.org/10.1088/2041-8205/789/1/L8)
- Chen, H. S.-L., & Shawhan, S. D. 1978, *SoPh*, 57, 205, doi: [10.1007/BF00152055](https://doi.org/10.1007/BF00152055)
- Chen, X., Kontar, E. P., Chrysaphi, N., et al. 2020, *ApJ*, 905, 43, doi: [10.3847/1538-4357/abc24e](https://doi.org/10.3847/1538-4357/abc24e)
- Chen, X., Kontar, E. P., Clarkson, D. L., & Chrysaphi, N. 2023, *MNRAS*, 520, 3117, doi: [10.1093/mnras/stad325](https://doi.org/10.1093/mnras/stad325)
- Chhetri, R., Morgan, J., Ekers, R. D., et al. 2018, *MNRAS*, 474, 4937, doi: [10.1093/mnras/stx2864](https://doi.org/10.1093/mnras/stx2864)
- Chiba, S., Imamura, T., Tokumaru, M., et al. 2022, *SoPh*, 297, 34, doi: [10.1007/s11207-022-01968-9](https://doi.org/10.1007/s11207-022-01968-9)
- Chrysaphi, N., Kontar, E. P., Holman, G. D., & Temmer, M. 2018, *ApJ*, 868, 79, doi: [10.3847/1538-4357/aae9e5](https://doi.org/10.3847/1538-4357/aae9e5)
- Chrysaphi, N., Reid, H. A. S., & Kontar, E. P. 2020, *ApJ*, 893, 115, doi: [10.3847/1538-4357/ab80c1](https://doi.org/10.3847/1538-4357/ab80c1)
- Clarkson, D. L., Kontar, E. P., Gordovskyy, M., Chrysaphi, N., & Vilmer, N. 2021, *ApJL*, 917, L32, doi: [10.3847/2041-8213/ac1a7d](https://doi.org/10.3847/2041-8213/ac1a7d)
- Clarkson, D. L., Kontar, E. P., Vilmer, N., et al. 2023, *ApJ*, 946, 33, doi: [10.3847/1538-4357/acbd3f](https://doi.org/10.3847/1538-4357/acbd3f)
- Cohen, M. H., Gundermann, E. J., Hardebeck, H. E., & Sharp, L. E. 1967, *ApJ*, 147, 449, doi: [10.1086/149028](https://doi.org/10.1086/149028)
- Coles, W. A. 1978, *SSRv*, 21, 411, doi: [10.1007/BF00173067](https://doi.org/10.1007/BF00173067)
- Coles, W. A., & Harmon, J. K. 1989, *ApJ*, 337, 1023, doi: [10.1086/167173](https://doi.org/10.1086/167173)
- Dennison, P. A., & Blesing, R. G. 1972, *Proceedings of the Astronomical Society of Australia*, 2, 86, doi: [10.1017/S1323358000012959](https://doi.org/10.1017/S1323358000012959)
- Dulk, G. A., & McLean, D. J. 1978, *SoPh*, 57, 279, doi: [10.1007/BF00160102](https://doi.org/10.1007/BF00160102)
- Dulk, G. A., & Suzuki, S. 1980, *A&A*, 88, 203
- Elgaroy, O., & Lyngstad, E. 1972, *A&A*, 16, 1
- Fainberg, J., & Stone, R. G. 1974, *SSRv*, 16, 145, doi: [10.1007/BF00240885](https://doi.org/10.1007/BF00240885)
- Fokker, A. D. 1965, *BAN*, 18, 111
- Fox, N. J., Velli, M. C., Bale, S. D., et al. 2016, *SSRv*, 204, 7, doi: [10.1007/s11214-015-0211-6](https://doi.org/10.1007/s11214-015-0211-6)
- Fredricks, R. W., & Coroniti, F. V. 1976, *J. Geophys. Res.*, 81, 5591, doi: [10.1029/JA081i031p05591](https://doi.org/10.1029/JA081i031p05591)
- Ginzburg, V. L., & Zhelezniakov, V. V. 1958, *Soviet Ast.*, 2, 653
- Gopalswamy, N., Nitta, N., Akiyama, S., Mäkelä, P., & Yashiro, S. 2012, *ApJ*, 744, 72, doi: [10.1088/0004-637X/744/1/72](https://doi.org/10.1088/0004-637X/744/1/72)
- Gordovskyy, M., Kontar, E., Browning, P., & Kuznetsov, A. 2019, *ApJ*, 873, 48, doi: [10.3847/1538-4357/ab03d8](https://doi.org/10.3847/1538-4357/ab03d8)
- Gorgolewski, S., Hanasz, J., Iwaniszewski, H., & Turło, Z. 1962, *AcA*, 12, 251
- Gurnett, D. A., Baumbach, M. M., & Rosenbauer, H. 1978, *J. Geophys. Res.*, 83, 616, doi: [10.1029/JA083iA02p00616](https://doi.org/10.1029/JA083iA02p00616)
- Harries, J. R., Blesing, R. G., & Dennison, P. A. 1970, *PASA*, 1, 319, doi: [10.1017/S1323358000012091](https://doi.org/10.1017/S1323358000012091)
- Hellinger, P., Matteini, L., Štverák, Š., Trávníček, P. M., & Marsch, E. 2011, *Journal of Geophysical Research (Space Physics)*, 116, A09105, doi: [10.1029/2011JA016674](https://doi.org/10.1029/2011JA016674)
- Hewish, A. 1958, *MNRAS*, 118, 534, doi: [10.1093/mnras/118.6.534](https://doi.org/10.1093/mnras/118.6.534)
- Hewish, A., Scott, P. F., & Wills, D. 1964, *Nature*, 203, 1214, doi: [10.1038/2031214a0](https://doi.org/10.1038/2031214a0)
- Hewish, A., & Wyndham, J. D. 1963, *MNRAS*, 126, 469, doi: [10.1093/mnras/126.5.469](https://doi.org/10.1093/mnras/126.5.469)
- Högbom, J. A. 1960, *MNRAS*, 120, 530, doi: [10.1093/mnras/120.6.530](https://doi.org/10.1093/mnras/120.6.530)

- Holman, G. D., Aschwanden, M. J., Aurass, H., et al. 2011, *SSRv*, 159, 107, doi: [10.1007/s11214-010-9680-9](https://doi.org/10.1007/s11214-010-9680-9)
- Kellogg, P. J., & Horbury, T. S. 2005, *Annales Geophysicae*, 23, 3765, doi: [10.5194/angeo-23-3765-2005](https://doi.org/10.5194/angeo-23-3765-2005)
- Kolmogorov, A. 1941, *Akademiia Nauk SSSR Doklady*, 30, 301
- Kolotkov, D. Y., Nakariakov, V. M., & Kontar, E. P. 2018, *ApJ*, 861, 33, doi: [10.3847/1538-4357/aac77e](https://doi.org/10.3847/1538-4357/aac77e)
- Kontar, E. P. 2001, *SoPh*, 202, 131, doi: [10.1023/A:1011894830942](https://doi.org/10.1023/A:1011894830942)
- Kontar, E. P., Yu, S., Kuznetsov, A. A., et al. 2017, *Nature Communications*, 8, 1515, doi: [10.1038/s41467-017-01307-8](https://doi.org/10.1038/s41467-017-01307-8)
- Kontar, E. P., Chen, X., Chrysaphi, N., et al. 2019, *ApJ*, 884, 122, doi: [10.3847/1538-4357/ab40bb](https://doi.org/10.3847/1538-4357/ab40bb)
- Krupar, V., Maksimovic, M., Santolik, O., Cecconi, B., & Kruparova, O. 2014, *SoPh*, 289, 4633, doi: [10.1007/s11207-014-0601-z](https://doi.org/10.1007/s11207-014-0601-z)
- Krupar, V., Maksimovic, M., Kontar, E. P., et al. 2018, *ApJ*, 857, 82, doi: [10.3847/1538-4357/aab60f](https://doi.org/10.3847/1538-4357/aab60f)
- Krupar, V., Szabo, A., Maksimovic, M., et al. 2020, *ApJS*, 246, 57, doi: [10.3847/1538-4365/ab65bd](https://doi.org/10.3847/1538-4365/ab65bd)
- Kuznetsov, A. A., Chrysaphi, N., Kontar, E. P., & Motorina, G. 2020, *ApJ*, 898, 94, doi: [10.3847/1538-4357/aba04a](https://doi.org/10.3847/1538-4357/aba04a)
- Kuznetsov, A. A., & Kontar, E. P. 2019, *A&A*, 631, L7, doi: [10.1051/0004-6361/201936447](https://doi.org/10.1051/0004-6361/201936447)
- Leamon, R. J., Smith, C. W., Ness, N. F., Matthaeus, W. H., & Wong, H. K. 1998, *J. Geophys. Res.*, 103, 4775, doi: [10.1029/97JA03394](https://doi.org/10.1029/97JA03394)
- Leblanc, Y., Dulk, G. A., & Bougeret, J.-L. 1998, *SoPh*, 183, 165, doi: [10.1023/A:1005049730506](https://doi.org/10.1023/A:1005049730506)
- Lee, L. C., & Jokipii, J. R. 1975, *ApJ*, 196, 695, doi: [10.1086/153458](https://doi.org/10.1086/153458)
- Lin, H., Penn, M. J., & Tomczyk, S. 2000, *ApJL*, 541, L83, doi: [10.1086/312900](https://doi.org/10.1086/312900)
- Lotz, S., Nel, A. E., Wicks, R. T., et al. 2023, *ApJ*, 942, 93, doi: [10.3847/1538-4357/aca903](https://doi.org/10.3847/1538-4357/aca903)
- Machin, K. E., & Smith, F. G. 1952, *Nature*, 170, 319, doi: [10.1038/170319b0](https://doi.org/10.1038/170319b0)
- Maguire, C. A., Carley, E. P., Zucca, P., Vilmer, N., & Gallagher, P. T. 2021, *ApJ*, 909, 2, doi: [10.3847/1538-4357/abda51](https://doi.org/10.3847/1538-4357/abda51)
- Mancuso, S., & Spangler, S. R. 2000, *ApJ*, 539, 480, doi: [10.1086/309205](https://doi.org/10.1086/309205)
- Mann, G., Jansen, F., MacDowall, R. J., Kaiser, M. L., & Stone, R. G. 1999, *A&A*, 348, 614
- Manoharan, P. K. 1993a, *SoPh*, 148, 153, doi: [10.1007/BF00675541](https://doi.org/10.1007/BF00675541)
- . 1993b, *Bulletin of the Astronomical Society of India*, 21, 383
- Marsch, E., & Tu, C. Y. 1990, *J. Geophys. Res.*, 95, 11945, doi: [10.1029/JA095iA08p11945](https://doi.org/10.1029/JA095iA08p11945)
- McCauley, P. I., Cairns, I. H., & Morgan, J. 2018, *SoPh*, 293, 132, doi: [10.1007/s11207-018-1353-y](https://doi.org/10.1007/s11207-018-1353-y)
- McKim Malville, J., Aller, H. D., & Jensen, C. J. 1967, *ApJ*, 147, 711, doi: [10.1086/149048](https://doi.org/10.1086/149048)
- Miyamoto, M., Imamura, T., Tokumaru, M., et al. 2014, *ApJ*, 797, 51, doi: [10.1088/0004-637X/797/1/51](https://doi.org/10.1088/0004-637X/797/1/51)
- Mohan, A. 2021, *A&A*, 655, A77, doi: [10.1051/0004-6361/202142029](https://doi.org/10.1051/0004-6361/202142029)
- Morimoto, M., & Kai, K. 1961, *PASJ*, 13, 294
- Murphy, P. C., Carley, E. P., Ryan, A. M., Zucca, P., & Gallagher, P. T. 2021, *A&A*, 645, A11, doi: [10.1051/0004-6361/202038518](https://doi.org/10.1051/0004-6361/202038518)
- Musset, S., Maksimovic, M., Kontar, E., et al. 2021, *A&A*, 656, A34, doi: [10.1051/0004-6361/202140998](https://doi.org/10.1051/0004-6361/202140998)
- Narayan, R., Anantharamaiah, K. R., & Cornwell, T. J. 1989, *MNRAS*, 241, 403, doi: [10.1093/mnras/241.3.403](https://doi.org/10.1093/mnras/241.3.403)
- Newkirk, Jr., G. 1961, *ApJ*, 133, 983, doi: [10.1086/147104](https://doi.org/10.1086/147104)
- Parker, E. N. 1958, *ApJ*, 128, 664, doi: [10.1086/146579](https://doi.org/10.1086/146579)
- Patzold, M., Bird, M. K., Volland, H., et al. 1987, *SoPh*, 109, 91, doi: [10.1007/BF00167401](https://doi.org/10.1007/BF00167401)
- Pulupa, M., Bale, S. D., Badman, S. T., et al. 2020, *ApJS*, 246, 49, doi: [10.3847/1538-4365/ab5dc0](https://doi.org/10.3847/1538-4365/ab5dc0)
- Ramesh, R., Kathiravan, C., & Sastry, C. V. 2010, *ApJ*, 711, 1029, doi: [10.1088/0004-637X/711/2/1029](https://doi.org/10.1088/0004-637X/711/2/1029)
- Reid, H. A. S., & Kontar, E. P. 2018, *A&A*, 614, A69, doi: [10.1051/0004-6361/201732298](https://doi.org/10.1051/0004-6361/201732298)
- . 2021, *Nature Astronomy*, 5, 796, doi: [10.1038/s41550-021-01370-8](https://doi.org/10.1038/s41550-021-01370-8)
- Reid, H. A. S., Vilmer, N., & Kontar, E. P. 2014, *A&A*, 567, A85, doi: [10.1051/0004-6361/201321973](https://doi.org/10.1051/0004-6361/201321973)
- Reiner, M. J., Fainberg, J., Kaiser, M. L., & Stone, R. G. 1998, *J. Geophys. Res.*, 103, 1923, doi: [10.1029/97JA02646](https://doi.org/10.1029/97JA02646)
- Reiner, M. J., Goetz, K., Fainberg, J., et al. 2009, *SoPh*, 259, 255, doi: [10.1007/s11207-009-9404-z](https://doi.org/10.1007/s11207-009-9404-z)
- Rickett, B. J. 1977, *ARA&A*, 15, 479, doi: [10.1146/annurev.aa.15.090177.002403](https://doi.org/10.1146/annurev.aa.15.090177.002403)
- Riddle, A. C. 1974, *SoPh*, 35, 153, doi: [10.1007/BF00156964](https://doi.org/10.1007/BF00156964)
- Roberts, O. W., Narita, Y., & Escoubet, C. P. 2018, *Annales Geophysicae*, 36, 527, doi: [10.5194/angeo-36-527-2018](https://doi.org/10.5194/angeo-36-527-2018)
- Rosenberg, R. L., Kivelson, M. G., Coleman, P. J., J., & Smith, E. J. 1978, *J. Geophys. Res.*, 83, 4165, doi: [10.1029/JA083iA09p04165](https://doi.org/10.1029/JA083iA09p04165)
- Saint-Hilaire, P., Vilmer, N., & Kerdran, A. 2013, *ApJ*, 762, 60, doi: [10.1088/0004-637X/762/1/60](https://doi.org/10.1088/0004-637X/762/1/60)

- Saito, K., Poland, A. I., & Munro, R. H. 1977, *SoPh*, 55, 121, doi: [10.1007/BF00150879](https://doi.org/10.1007/BF00150879)
- Sakurai, T., & Spangler, S. R. 1994, *ApJ*, 434, 773, doi: [10.1086/174780](https://doi.org/10.1086/174780)
- Sasikumar Raja, K., Ingale, M., Ramesh, R., et al. 2016, *Journal of Geophysical Research (Space Physics)*, 121, 11, doi: [10.1002/2016JA023254](https://doi.org/10.1002/2016JA023254)
- Sasikumar Raja, K., Subramanian, P., Ramesh, R., Vourlidas, A., & Ingale, M. 2017, *ApJ*, 850, 129, doi: [10.3847/1538-4357/aa94cd](https://doi.org/10.3847/1538-4357/aa94cd)
- Sasikumar Raja, K., Maksimovic, M., Kontar, E. P., et al. 2022, *ApJ*, 924, 58, doi: [10.3847/1538-4357/ac34ed](https://doi.org/10.3847/1538-4357/ac34ed)
- Scott, S. L., Coles, W. A., & Bourgois, G. 1983, *A&A*, 123, 207
- Shain, C. A., & Higgins, C. S. 1959, *Australian Journal of Physics*, 12, 357, doi: [10.1071/PH590357](https://doi.org/10.1071/PH590357)
- Sharma, R., & Oberoi, D. 2021, *ApJ*, 913, 153, doi: [10.3847/1538-4357/ac01df](https://doi.org/10.3847/1538-4357/ac01df)
- Sharykin, I. N., Kontar, E. P., & Kuznetsov, A. A. 2018, *SoPh*, 293, 115, doi: [10.1007/s11207-018-1333-2](https://doi.org/10.1007/s11207-018-1333-2)
- Shevchuk, N. V., Melnik, V. N., Poedts, S., et al. 2016, *SoPh*, 291, 211, doi: [10.1007/s11207-015-0799-4](https://doi.org/10.1007/s11207-015-0799-4)
- Sioulas, N., Huang, Z., Shi, C., et al. 2023, *ApJL*, 943, L8, doi: [10.3847/2041-8213/acaeff](https://doi.org/10.3847/2041-8213/acaeff)
- Sittler, Edward C., J., & Guhathakurta, M. 1999, *ApJ*, 523, 812, doi: [10.1086/307742](https://doi.org/10.1086/307742)
- Slee, O. B. 1959, *Australian Journal of Physics*, 12, 134, doi: [10.1071/PH590134](https://doi.org/10.1071/PH590134)
- . 1966, *Planet. Space Sci.*, 14, 255, doi: [10.1016/0032-0633\(66\)90125-5](https://doi.org/10.1016/0032-0633(66)90125-5)
- Spangler, S. R. 2002, *ApJ*, 576, 997, doi: [10.1086/341889](https://doi.org/10.1086/341889)
- . 2005, *SSRv*, 121, 189, doi: [10.1007/s11214-006-4719-7](https://doi.org/10.1007/s11214-006-4719-7)
- . 2020, *Research Notes of the American Astronomical Society*, 4, 147, doi: [10.3847/2515-5172/abb29a](https://doi.org/10.3847/2515-5172/abb29a)
- Spangler, S. R., Kavars, D. W., Kortenkamp, P. S., et al. 2002, *A&A*, 384, 654, doi: [10.1051/0004-6361:20020028](https://doi.org/10.1051/0004-6361:20020028)
- Steinberg, J. L. 1972, *A&A*, 18, 382
- Steinberg, J. L., Aubier-Giraud, M., Leblanc, Y., & Boisshot, A. 1971, *A&A*, 10, 362
- Steinberg, J. L., Hoang, S., & Dulk, G. A. 1985, *A&A*, 150, 205
- Stewart, R. T. 1972, *PASA*, 2, 100, doi: [10.1017/S1323358000013059](https://doi.org/10.1017/S1323358000013059)
- Suzuki, S., & Dulk, G. A. 1985, in *Solar Radiophysics: Studies of Emission from the Sun at Metre Wavelengths*, ed. D. J. McLean & N. R. Labrum (Cambridge University Press), 289–332
- Tasnim, S., Zank, G. P., Cairns, I. H., & Adhikari, L. 2022, *ApJ*, 928, 125, doi: [10.3847/1538-4357/ac5031](https://doi.org/10.3847/1538-4357/ac5031)
- Thejappa, G., & MacDowall, R. J. 2008, *ApJ*, 676, 1338, doi: [10.1086/528835](https://doi.org/10.1086/528835)
- Tyul'bashev, S. A., Chashei, I. V., & Kitaeva, M. A. 2023, *MNRAS*, doi: [10.1093/mnras/stad1461](https://doi.org/10.1093/mnras/stad1461)
- Villante, U., & Mariani, F. 1975, *Geophys. Res. Lett.*, 2, 73, doi: [10.1029/GL002i003p00073](https://doi.org/10.1029/GL002i003p00073)
- Šafránková, J., Němeček, Z., Němec, F., et al. 2015, *ApJ*, 803, 107, doi: [10.1088/0004-637X/803/2/107](https://doi.org/10.1088/0004-637X/803/2/107)
- Wild, J. P., Sheridan, K. V., & Trent, G. H. 1959, in *IAU Symposium, Vol. 9, URSI Symp. 1: Paris Symposium on Radio Astronomy*, ed. R. N. Bracewell, 176
- Woo, R. 1977, *Nature*, 266, 514, doi: [10.1038/266514a0](https://doi.org/10.1038/266514a0)
- . 1978, *ApJ*, 219, 727, doi: [10.1086/155831](https://doi.org/10.1086/155831)
- Woo, R., Yang, F. C., & Ishimaru, A. 1977, *ApJ*, 218, 557, doi: [10.1086/155710](https://doi.org/10.1086/155710)
- Yakovlev, O. I., Efimov, A. I., Razmanov, V. M., & Shtrykov, V. K. 1980, *Soviet Ast.*, 24, 454
- Yamauchi, Y., Tokumaru, M., Kojima, M., Manoharan, P. K., & Esser, R. 1998, *J. Geophys. Res.*, 103, 6571, doi: [10.1029/97JA03598](https://doi.org/10.1029/97JA03598)
- Yamauchi, Y., Tokumaru, M., Kojima, M., et al. 1996, in *American Institute of Physics Conference Series, Vol. 382, Proceedings of the eighth International solar wind Conference: Solar wind eight*, ed. D. Winterhalter, J. T. Gosling, S. R. Habbal, W. S. Kurth, & M. Neugebauer, 366–366, doi: [10.1063/1.51472](https://doi.org/10.1063/1.51472)
- Zank, G. P., Adhikari, L., Hunana, P., et al. 2017, *ApJ*, 835, 147, doi: [10.3847/1538-4357/835/2/147](https://doi.org/10.3847/1538-4357/835/2/147)
- Zhang, P., Yu, S., Kontar, E. P., & Wang, C. 2019, *ApJ*, 885, 140, doi: [10.3847/1538-4357/ab458f](https://doi.org/10.3847/1538-4357/ab458f)