

X-ray flares of the young planet host DS Tuc A.

I. Pillitteri¹, C. Argiroffi^{2,1}, A. Maggio¹, G. Micela¹, S. Benatti¹, F. Reale², S. Colombo¹, and S.J. Wolk³

¹ INAF-Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90134 Palermo, Italy
e-mail: ignazio.pillitteri@inaf.it

² Università degli Studi di Palermo, Piazza Marina 61, 90133 Palermo, Italy

³ Harvard-Smithsonian Center for Astrophysics, 02138 Garden St, Cambridge, MA

Received ; accepted

ABSTRACT

The discovery of planets around young stars has increased the study of the early phases of planetary formation and evolution. Stars are strong emitters at X-ray and UV wavelengths in the first billion of years and this strongly affects the evaporation, thermodynamics and chemistry of the atmospheres of young planets around them. In order to investigate these effects in young exoplanets we observed the 40 Myr old star DS Tuc A with XMM-Newton and recorded two X-ray bright flares, with the second event occurring about 12 ks after the first one. Their duration from the rise to the end of the decay was of about 8 – 10 ks in soft X-rays (0.3-10 keV). The flares were also recorded in the band 200–300 nm with the UVM2 filter of the Optical Monitor. The duration of the flares in UV was about 3 ks. The observed delay between the peak in the UV band and in X-rays is a probe of the heating phase followed by the evaporation and increase of density and emission measure of the flaring loop. The coronal plasma temperature at the two flare peaks reached 54–55 MK. The diagnostics based on temperatures and time scales of the flares applied to these two events allow us to infer a loop length of $5 - 7 \times 10^{10}$ cm, which is about the size of the stellar radius. We also infer values of electron density at the flare peaks of $2.3 - 6.5 \times 10^{11}$ cm⁻³, and a minimum magnetic field strength of order of 300–500 G needed to confine the plasma. The energy released during the flares was of order of $5 - 8 \times 10^{34}$ erg in the band 0.3 – 10 keV and $0.9 - 2.7 \times 10^{33}$ erg in the UV band (200-300 nm). We speculate that the flares were associated with Coronal Mass Ejections (CMEs) that hit the planet about 3.3 hr after the flares and dramatically increasing the rate of evaporation of the planet. From the RGS spectra we retrieved the emission measure distribution and the abundances of coronal metals during the quiescent and the flaring states. In agreement with what inferred from time resolved spectroscopy and EPIC spectra, also from the analysis of RGS spectra during the flares we infer a high electron density.

Key words. Stars: flare – Stars: coronae – Stars: activity – X-rays: stars

1. Introduction

X-rays are emitted by solar type stars during their early evolution (Vaiana et al. 1981; Pallavicini et al. 1981; Feigelson & Montmerle 1999) and are a powerful proxy for the structure of the stellar coronae and the magnetic fields that confine the hot coronal plasma. During the first hundred million years, solar type stars exhibit very high magnetic and coronal activity, as well as high stellar rotation rates (Pizzolato et al. 2003). The loss of angular momentum during the Main Sequence is accompanied by a decrease of the X-ray luminosity by a factor of 10^3 during the first billion of years (Pallavicini et al. 1981; Vaiana et al. 1981; Favata et al. 2003).

The interest for the study of the activity of stars at high energies has been renewed with the discovery of extrasolar planets. High energy photons and particles can affect planetary atmospheres and their habitability. Fluxes at high energies (X-rays and EUV, XUV) have a strong role in shaping and transforming the primary atmospheres of the planets and their evaporation. Because of the decrease of stellar activity, most of the effects due to XUV irradiation are expected to happen in the first Gyr (Penz & Micela 2008; Cecchi-Pestellini et al. 2009; Kubyskhina et al. 2018). Collectively, the action of the magnetized stellar winds and the coronal mass ejections (CMEs) can make the planets lose a significant fraction of their atmospheres and change their compositions through photo-chemical processes (see Drake et al. 2013 and Kay et al. 2019 and references therein). However, the

effects of the sudden and violent release of energy during flares and CMEs onto the planets are still poorly understood due to the lack of proper monitoring of the stellar coronae required in order to build a reliable distribution of flaring rates vs. age, and an adequate time-resolved modeling of the dynamics and chemistry of the planetary gaseous envelopes. In this context, the study of young stars hosting planets, like DS Tuc, in X-ray and UV bands is of high relevance.

DS Tuc is composed by a pair of two young stars at a distance of 44.1 pc from the Sun, with an age of ~ 40 Myr and spectral types G6 (component A and host of the planet) and K3 (component B). The planet around DS Tuc A was independently discovered in NASA TESS observations (TOI-200) by Benatti et al. (2019) and Newton et al. (2019). DS Tuc Ab has a radius of 0.5 R_J, a mass upper limit of 14.4 M_⊕ (Benatti et al. 2021) and orbits its host star in 8.14 days. These characteristics suggest a very low density for the planet and that it is prone to strong atmospheric evaporation. This is one of few very young planets discovered so far. Its properties can serve to test models of dynamical and atmospheric evolution. For the purpose of investigating its coronal emission, DS Tuc has been observed with XMM-Newton in two different observations. In Benatti et al. (2021) we have presented the analysis and the global properties of the coronae of the system of DS Tuc and related the X-ray emission to the model of atmospheric evaporation during the Main Sequence.

In the optical band TESS has observed DS Tuc A in six sectors allowing Colombo et al. (2022) to perform a reconstruction

of the distribution of the optical flares and their main properties. They also inferred the energies of the X-ray counterparts of TESS flares through the scaling laws of Flaccomio et al. (2018) before X-rays observations were available. Here, we study two bright flares of DS Tuc A and discuss their impact on the atmosphere of DS Tuc Ab.

The paper is structured as follows: Sect. 2 describes the observations and the data analysis, Sect. 3 presents our results and Sect. 4 contains our discussion and conclusions.

2. XMM-Newton observations and data analysis

DS Tuc has been observed two times in X-rays with *XMM-Newton* during observations ObsId=0863400901 (P.I. S. Wolk) and 0864340101 (P.I. A. Maggio) with exposure times of about 40 ks and 30 ks, respectively. From the first observation (Benatti et al. 2021) we measured X-ray fluxes and luminosities of the pair of stars of DS Tuc A and B, separated by $\sim 5.3''$ and partially resolved in MOS detectors. The A component was 1.3 times more luminous in X-rays than DS Tuc B. We also inferred a mean coronal temperature and emission measure of both stars. Some degree of variability in form of flares was observed in DS Tuc B while DS Tuc A was quieter. From these we estimated the time scale of the evaporation of DS Tuc Ab and the evolution of its mass and radius in the next 5 Gyr.

The second observation of DS Tuc with *XMM-Newton* was performed on April 11th 2021 (obsId 0864340101)¹, again with EPIC as main instrument. We adopted the *Medium* filter and the *SmallWindow* mode for both MOS and *pn*. In addition, we acquired simultaneous high resolution spectra with RGS and a light curve in the band 200 – 300 nm with the Optical Monitor (OM) using the UVM2 filter in Fast mode. For the OM we obtained five continuous exposures of 4400 s and one exposure of 3880 s, for each exposure a light curve with a time binning of 10 s was obtained.

The ODFs files downloaded from the *XMM-Newton* archive² were reduced with SAS ver 18.0.0, to obtain event tables for EPIC instruments, light curves for OM and spectra for the two RGS high-resolution spectrometers. We selected the EPIC events for having energies in the band 0.3 – 10.0 keV and FLAG == 0 and PATTERN <= 12 as prescribed by the SAS reduction guide. We checked that the overall background was low during the observation, thus the full exposure was retained for further analysis. A visual inspection of the *pn* and MOS images revealed that DS Tuc A underwent strong flaring activity and its flux dominated the emission of the nearby DS Tuc B. Fig. 1 shows two MOS1 images relative to two different time intervals and referring to the quiescent and flare peak intervals of DS Tuc A.

Spectra and light curves of DS Tuc A were obtained from events in a circular region of radius $75''$ centered on the X-ray image centroid (see Fig. 1). Spectra of the background were accumulated from an offset circular region of radius $45''$. Given the brightness of DS Tuc A in this observation we checked for any pile up that could have affected the spectra of MOS and *pn*. We generated the pattern of single, double, triple, and quadruple events as a function of the energy as prescribed by the SAS guide. The diagnostic modeling of the curve did not show evidence of significant pile up so we retained the entire list of events in the region.

¹ A detailed log of the observation is available at: https://xmmweb.esac.esa.int/cgi-bin/xmmobs/public/obs_view.tcl?search_obs_id=0864340101

² <https://nxs.a.esac.esa.int/nxs-a-web/#home>




Fig. 1. Top panel: MOS 2 image of DS Tuc with the regions used for accumulating source and background events. The scale is logarithmic to enhance the background level otherwise not visible on a linear scale. The background region was offset to one of the corner of the chip to avoid the diffraction spikes from the central source. Bottom left: an image of MOS 1 during the first 6 ks of observation and corresponding to interval 1 in Fig. 2. We marked the SIMBAD positions (J2000) of DS Tuc A and B and the vectors of proper motions. Bottom right: image during interval 3 corresponding to the peak of the first flare. In both images the scale limits and color maps are identical, the binning is 8 pixels corresponding to $\sim 0.4''$ and a smoothing with a Gaussian with $\sigma = 1.5$ pixels has been applied to images. We added contour levels with the same ranges for marking the source centroid.

The high count statistics recorded in the EPIC instruments allowed us to perform time resolved spectroscopy of the two flares. To this purpose we accumulated the spectra in 13 time intervals that define the initial quiescent state, the flare rises, peaks and decays (see Fig. 2). For each time interval the spectra of the source and the background were created with SAS, along with the corresponding response matrices and effective area ARF files.

For the OM we obtained five exposures of 4400 s and one of 3880 s. In each exposure, a light curve with time binning of 10 s was accumulated with the standard SAS task *omfchain*, and the result is displayed in Fig. 2.

For extracting the RGS spectra we used the SAS task *rgsproc* and *rgscombine* with custom good time intervals defined as in-




Fig. 2. Light curves of MOS 1 (red curve, bin size 150 s) and OM in UVM2 filter (blue curve, bin size 10 s). OM rate is scaled down by a factor 30 to allow a direct comparison of the rise and decay times of both instruments during the flares. The time intervals used for time resolved spectroscopy (see Sect. 3.2) are indicated by the alternating light/dark grey areas and numbered on top of each panel.




Fig. 3. Spectra of MOS 1 relative to the time intervals 1 (black line and crosses) and 3 (red crosses). The best fit model to the spectrum of the plasma during interval 1 with its 3 component is shown along with the χ residuals in the bottom panel.

interval nr. 1 for quiescent and 2 to 13 for flares. We summed the RGS spectrum of the first *XMM-Newton* observation (ObsId 0863400901) to the spectrum of the quiescent interval nr. 1, since both are representative of a quiescent phase or at least with low level of variability. Furthermore, we added RGS 1 and RGS 2 spectra of the quiescent and flaring phases for achieving a better count statistics.

3. Results

We discuss separately the quiescent emission and the flaring emission observed in the present observation.

3.1. Quiescent phase

The quiescent interval before the first flare was about 6.5 ks with an average rate of 0.9 ct/s in MOS1, this interval served to estimate the coronal plasma properties before the ignition. We used

a sum of 3 APEC thermal components plus a global absorption to model the EPIC spectra and obtain a statistically satisfactory description of the spectrum. Table 1 lists the best fit parameters and Fig. 3 shows the spectrum during time interval 1 and 3 for comparison. We performed the best fit procedure to each EPIC spectrum and to them jointly obtaining very similar results.

The main thermal component of the quiescent spectrum is $\log(T/K) \sim 7.02$, with the addition of a soft component $\log T/K \sim 0.54$ and a hotter component, much less constrained, $\log T/K \geq 7.4$. By setting a fixed very low value to the equivalent column of gas absorption ($N_H = 8 \times 10^{18} \text{ cm}^{-2}$) we can constrain the hot component at around $\log T \sim 7.4$ (90% confidence range: 7.31–7.58) while the other parameters remain unchanged. However the very low gas absorption is not consistent with the values of visual extinction and reddening.

The temperatures are consistent with the values determined from the analysis of spectra acquired during the first *XMM-Newton* observation (Benatti et al. 2021). The average flux of DS Tuc A in interval 1 is about 1.9 times the flux observed during the first *XMM-Newton* observation. However, some degree of contamination from DS Tuc B in this estimate has to be accounted for in the analysis of this first time interval. Visual inspection of the image relative to the events of the quiescent interval shows that DS Tuc A is still the most prominent source and DS Tuc B has a lower count statistics. The contribution to flux of DS Tuc A from its stellar companion DS Tuc B is of order of 40% during interval 1. After correcting for this contamination fraction, we infer that the quiescent flux of DS Tuc A in interval 1 was about 15% higher than the average X-ray flux measured during the first *XMM-Newton* observation.

3.2. Flares

We observed two prominent flares after the initial phase of quiescence that lasted for about 6.5 ks (Fig. 2) from the start of the *XMM-Newton* observation. The second flare occurred about 12 ks after the beginning of the first flare. In Fig. 2 we show a light curve of the OM with scaled count rate to allow a straightforward comparison of the timing of the flares. Each flare starts simultaneously in X-rays and UV, however in the UV band the flares peak earlier and decays quicker than in X-ray. When the flux reaches each flare peak in UV the rise of the flux in X-rays does flatten while continuing to increase. The peak in UV corresponds to the end of the heating phase during the flares while the loop continues to be filled with plasma evaporating from its feet and thus increasing the emission measure and the flux in X-rays (Namekata et al. 2017). This timing and shape of the flares in the near UV band and in X-rays is analog to the Neupert effect seen in the Sun (Neupert 1968). Neupert noticed that time-integrated microwave fluxes closely match the rising portions of soft X-ray (SXR) emission. The effect has been demonstrated since by others (e.g., Kahler et al. 1988), and was later dubbed as the Neupert effect by Hudson (1991). Estimating the time-integrated microwave fluxes from the non-thermal hard X-ray (HXR) emission, the Neupert effect has been generalized, where the time derivative of the SXR, i.e., $dF_{SXR}(t)/dt$, can be considered as a proxy for the HXR emission $F_{HXR}(t)$ (Dennis & Zarro 1993, Veronig et al. 2002). In our case, near UV flux measured with OM is a proxy of the dF_{HXR}/dt , with HXR being the emission in 0.3–10 keV probed by EPIC.

In the UV we measured rise times of 180 ± 5 s and 70 ± 5 s for the first and second flare respectively. In X-rays we measured rise times of 900 ± 75 s and 750 ± 75 s, respectively. The delay between the peak of each flare in UV and X-rays amount to ~

Table 1. Best fit parameters of the 3T model for the quiescent spectrum observed during interval 1. We list the parameters obtained from the best fit to each EPIC spectrum and the joint fit of *pn* and MOS. Unabsorbed flux and luminosity are calculated in the band 0.3–10.0 keV.

Parameter	MOS1	MOS2	<i>pn</i>	Joint
N_{H} (10^{20})	1.5 (0–4.4)	1.55 (0–4.25)	2.08 (0.44–3.68)	2.3 (1.0–3.4)
$\log T_1$ (K)	6.54 (6.48–6.61)	6.67 (6.6–6.76)	6.57 (6.52–6.62)	6.59 (6.56–6.63)
$\log EM1$ (cm^{-3})	52.89 (52.69–53.16)	52.95 (52.8–53.08)	52.99 (52.83–53.12)	53.02 (52.91–53.10)
$\log T_2$ (K)	7.03 (6.99–7.05)	7.06 (7.04–7.07)	7.01 (7–7.03)	7.03 (7.02–7.04)
$\log EM2$ (cm^{-3})	52.95 (52.73–53.17)	53.12 (53.08–53.19)	53.13 (52.98–53.24)	53.14 (53.02–53.20)
$\log T_3$ (K)	7.42 (>7.25)	>7.64	7.58 (>7.34)	7.87 (> 7.53)
$\log EM3$ (cm^{-3})	52.51 (51.97–52.73)	51.97 (51.67–52.15)	52.32 (51.97–52.6)	52.08 (> 51.91)
Z/Z_{\odot}	0.22 (0.13–0.34)	0.14 (0.11–0.17)	0.13 (0.0984–0.1804)	0.13 (0.11–0.16)
χ^2	52.05	55.74	60.37	190.82
D.o.F.	47	48	48	159
$P(\chi^2 > \chi_0^2)$	0.28	0.21	0.11	0.04
$\log f_{\text{X}}$ ($\text{erg s}^{-1} \text{cm}^{-2}$)	-11.11	-11.13	-11.1	-11.09
$\log L_{\text{X}}$ (erg s^{-1})	30.26	30.23	30.27	30.27

680 s and ~ 930 s, respectively. In a linear-log plane the decay phase of the flares appear first linear meaning that the first part of the decay can be modeled with an exponential relationship between count rate and time, then it does flatten deviating from a pure exponential decay. With a linear best fit modeling to the first part of the decay of both flares in the linear-log plane we estimated e-folding times of 3.55 ks and 2.31 ks for flares 1 and 2, respectively. With similar procedure we determined that in UV the flare decays are shorter than in X-rays and of order of 390 s and 375 s for flares 1 and 2, respectively.

The evolution of the plasma temperature and its emission measure can provide diagnostics about the size of the magnetic loop that hosted the flare, the presence of prolonged heating and the minimum magnetic field required to confine the plasma (Reale 2007, 2014). In order to follow the evolution of the flares we divided the light curve during the flares in 12 intervals that identify the rise, the peak and the decay (resolved with 3 different segments each) of each flare as indicated in Fig. 2. For each interval we modeled the extracted spectra with a sum of a flaring component described by one APEC optically thin plasma thermal emission model added to the 3-T model that describes the quiescent emission (cf. Sect. 3.1). We left free to vary also the global absorption and the global abundances. For each time segment the best fit parameters of spectra of MOS 1, 2 and *pn* are listed in Table 2. We did not perform a joint best fit to the 3 EPIC spectra of each time interval because the diagnostics of the flares are calibrated for each instrument separately. The absorption is of order of 10^{20} cm^{-2} , in a few cases it is not well constrained but we found the 90% confidence level value always below $4 \times 10^{20} \text{ cm}^{-2}$. These values are in agreement with the estimates of optical extinction towards DS Tuc (cf. Benatti et al. 2021).

Fig. 4 shows the scatter plots of the flare temperatures vs. the emission measures (E.M.) for each EPIC instrument. Points 2 to 7 refer to the first flare, points 8 to 13 refer to the second flare (cf. Table 2). In the spectra of MOS 1 and *pn* the peak temperatures in both flares are detected in segments 2 and 8 for flare 1 and 2, respectively, and of order of $\log T/\text{K} \sim 7.7 - 7.75$. In MOS 2 for the second flare the peak temperature is detected in segment 9, however its value is consistent with that measured in segment 8 with the other two instruments. The peaks of E.M. are reached in segments 3 and 9 for MOS 1 and *pn* with values of $\log EM(\text{cm}^{-3}) \approx 54$. Owing to our choice of the time intervals, the peak of E.M. in MOS 2 is seen in segment 8 instead of 9, but still consistent with the values determined from MOS 1 and *pn*. We inferred a metallicity value of Z between 0.2 during the decay and 0.5 at the flare peaks.

The slopes between points 3 and 7, and 9 to 13 in the $\log T / EM$ plane, the temperatures at the peaks of the flares and the e-folding time of the decay of the flares can be used to infer the size of the loops that hosted the flares using the diagnostics of Reale (2007) and Reale (2014). Table 3 shows the slopes derived for each flare and EPIC instrument, the total loop lengths, the estimates of the electron densities and the minimum strength of the magnetic field required to constrain the plasma in the loop at the peak of both flares. We inferred a full loop length of order of the stellar radius ($5 - 7 \times 10^{10} \text{ cm}$). To estimate the electron density we first calculated the volume of the flaring loops assumed as tubes with a ratio between base radius and length of 0.1. Then we derived the electron densities from E.M., as $n_e = \sqrt{EM/V}$ (Maggio et al. 2000). Electron densities at the flare peaks are found of order of $2 - 6.5 \times 10^{11} \text{ cm}^{-3}$. We can also infer the minimum strength of the magnetic field needed to confine the plasma as (e.g. Maggio et al. 2000)

$$B = \sqrt{16\pi k_B n_{e,\text{peak}} T_{\text{peak}}}$$

where k_B is the Boltzmann's constant, $n_{e,\text{peak}}$ and T_{peak} are the density and the temperature at the flare peak. We estimated a minimum magnetic field between 300 G and 500 G.

3.3. RGS lines

We analyzed the first-order RGS spectra of both *XMM-Newton* observations to infer the emission measure distribution (EMD), abundances, and electron density of the X-ray emitting plasma during the quiescent and flaring phases. First order RGS spectra cover the range $\approx 4 - 38 \text{ \AA}$ with a resolution 0.06 \AA FWHM. The exposures of the quiescent and flaring intervals are 45.53 ks and 22.15 ks long, respectively. The quiescent spectrum was accumulated summing up the events collected during the whole first *XMM-Newton* observation and the first 6.2 ks of the present observation. For the flaring phase we considered the remaining 22.2 ks encompassing the two flares altogether. The RGS spectra of the first order are made of 11200 and 20700 net counts, respectively. For the analysis of the RGS data we used the IDL package PINTofALE, (Kashyap & Drake 2000) and the XSPEC v12.11b software (Arnaud 1996). For the line emissivities we adopted the CHIANTI atomic database (v7.13, Dere et al. 1997).

We identified the strongest emission lines and measured their fluxes by fitting the RGS spectra in small wavelength intervals of $\sim 1.0 \text{ \AA}$ width. The width of each wavelength interval is set in order to include, and hence fit simultaneously, blended lines.

Table 2. Best fit parameters from time resolved spectroscopy of EPIC spectra. For each time interval (column 1) are reported the column equivalent absorption (N_{H}), the temperature (kT), the emission measure (E.M.) the global abundance scaling factor (Z/Z_{\odot}), the unabsorbed flux and luminosity in the 0.3-10 keV band, the χ^2 , the degrees of freedom and the probability related to the χ^2 statistics. The parameter ranges refer to the 90% confidence intervals.

Interval	N_{H} 10^{20} cm^{-2}	log (T/K)	log E.M. cm^{-3}	Z/Z_{\odot}	log f_{X} $\text{erg s}^{-1} \text{ cm}^{-2}$	log L_{X} erg s^{-1}	χ^2	d.o.f.	Prob
MOS1									
2	1.31 (0.17–2.55)	7.75 (7.67–7.83)	53.6 (53.57–53.63)	0.29 (0.25–0.34)	-10.45	30.92	52.05	47	0.28
3	0.71 (0.06–1.39)	7.62 (7.59–7.65)	53.96 (53.94–53.98)	0.56 (0.5–0.62)	-10.13	31.24	55.74	48	0.21
4	0 (0–0.58)	7.51 (7.47–7.54)	53.84 (53.81–53.87)	0.47 (0.41–0.53)	-10.29	31.08	60.37	48	0.109
5	0.45 (0–1.13)	7.51 (7.48–7.55)	53.58 (53.56–53.61)	0.49 (0.45–0.54)	-10.45	30.91	52.05	47	0.28
6	0.61 (0–1.32)	7.41 (7.37–7.46)	53.34 (53.31–53.37)	0.35 (0.31–0.38)	-10.68	30.69	55.74	48	0.21
7	0 (0–0.44)	7.36 (7.32–7.42)	53.17 (53.13–53.22)	0.3 (0.27–0.33)	-10.79	30.57	60.37	48	0.109
8	2.22 (1.02–3.52)	7.73 (7.67–7.81)	53.65 (53.62–53.68)	0.45 (0.39–0.52)	-10.37	31	52.05	47	0.28
9	1.67 (0.68–2.72)	7.66 (7.61–7.71)	53.93 (53.9–53.96)	0.49 (0.41–0.58)	-10.16	31.21	55.74	48	0.21
10	0.27 (0–1.49)	7.61 (7.56–7.67)	53.77 (53.74–53.8)	0.51 (0.43–0.59)	-10.29	31.07	60.37	48	0.109
11	0.83 (0.05–1.66)	7.53 (7.48–7.57)	53.6 (53.57–53.63)	0.44 (0.4–0.49)	-10.46	30.91	52.05	47	0.28
12	0.15 (0–0.92)	7.37 (7.32–7.43)	53.15 (53.1–53.19)	0.31 (0.28–0.34)	-10.79	30.57	55.74	48	0.21
13	1.28 (0.11–2.52)	7.2 (7.09–7.41)	52.22 (51.82–52.46)	0.25 (0.22–0.29)	-11.04	30.32	60.37	48	0.109
MOS2									
2	0.65 (0–1.96)	7.65 (7.58–7.73)	53.64 (53.59–53.66)	0.21 (0.17–0.26)	-10.48	30.88	52.05	47	0.28
3	2.04 (1.33–2.78)	7.61 (7.58–7.64)	53.96 (53.94–53.98)	0.54 (0.48–0.6)	-10.14	31.23	55.74	48	0.21
4	0.49 (0–1.41)	7.56 (7.51–7.6)	53.81 (53.78–53.84)	0.51 (0.44–0.58)	-10.28	31.09	60.37	48	0.109
5	1.06 (0.32–1.84)	7.5 (7.46–7.54)	53.57 (53.54–53.59)	0.46 (0.42–0.5)	-10.48	30.89	52.05	47	0.28
6	1.28 (0.54–2.06)	7.38 (7.34–7.42)	53.35 (53.31–53.38)	0.33 (0.3–0.37)	-10.69	30.68	55.74	48	0.21
7	0 (0–0.78)	7.38 (7.34–7.44)	53.16 (53.11–53.2)	0.29 (0.27–0.32)	-10.8	30.57	60.37	48	0.109
8	0.34 (0–1.57)	7.66 (7.59–7.73)	53.69 (53.65–53.72)	0.36 (0.3–0.42)	-10.39	30.98	52.05	47	0.28
9	1.23 (0.14–2.4)	7.73 (7.67–7.79)	53.88 (53.85–53.91)	0.51 (0.43–0.59)	-10.18	31.19	55.74	48	0.21
10	1.19 (0–2.48)	7.57 (7.52–7.63)	53.8 (53.77–53.84)	0.4 (0.32–0.49)	-10.31	31.06	60.37	48	0.109
11	0.5 (0–1.33)	7.53 (7.49–7.57)	53.59 (53.56–53.62)	0.4 (0.35–0.44)	-10.47	30.9	52.05	47	0.28
12	1.14 (0.27–2.08)	7.35 (7.29–7.41)	53.13 (53.07–53.18)	0.3 (0.27–0.34)	-10.81	30.55	55.74	48	0.21
13	2.58 (1.34–4.02)	7.09 (7.02–7.24)	52.45 (52.08–52.73)	0.25 (0.19–0.3)	-11.03	30.34	60.37	48	0.109
pn									
2	0.24 (0–0.81)	7.7 (7.66–7.75)	53.72 (53.7–53.74)	0.18 (0.16–0.2)	-10.37	30.99	52.05	47	0.28
3	0.16 (0–0.52)	7.64 (7.62–7.67)	54.05 (54.04–54.07)	0.38 (0.36–0.41)	-10.06	31.3	55.74	48	0.21
4	0.62 (0.15–1.08)	7.49 (7.47–7.52)	53.92 (53.9–53.94)	0.35 (0.32–0.37)	-10.23	31.14	60.37	48	0.109
5	0.56 (0.22–0.93)	7.45 (7.43–7.48)	53.66 (53.64–53.68)	0.3 (0.29–0.32)	-10.45	30.92	52.05	47	0.28
6	1.65 (1.28–2.05)	7.38 (7.35–7.41)	53.42 (53.4–53.44)	0.22 (0.21–0.23)	-10.65	30.71	55.74	48	0.21
7	1.54 (1.12–1.98)	7.34 (7.3–7.38)	53.28 (53.25–53.31)	0.19 (0.18–0.2)	-10.75	30.61	60.37	48	0.109
8	0.98 (0.43–1.56)	7.71 (7.66–7.75)	53.79 (53.77–53.81)	0.25 (0.23–0.27)	-10.3	31.07	52.05	47	0.28
9	0.08 (0–0.6)	7.67 (7.64–7.7)	54.01 (53.99–54.02)	0.31 (0.28–0.35)	-10.11	31.25	55.74	48	0.21
10	0.54 (0–1.14)	7.56 (7.52–7.6)	53.86 (53.84–53.89)	0.31 (0.28–0.34)	-10.26	31.1	60.37	48	0.109
11	0.25 (0–0.66)	7.52 (7.49–7.55)	53.67 (53.65–53.69)	0.26 (0.24–0.28)	-10.44	30.93	52.05	47	0.28
12	1.13 (0.72–1.56)	7.31 (7.28–7.36)	53.24 (53.21–53.28)	0.19 (0.17–0.2)	-10.78	30.59	55.74	48	0.21
13	1.28 (0.72–1.86)	7.14 (7.05–7.23)	52.58 (52.43–52.76)	0.13 (0.11–0.15)	-11.02	30.34	60.37	48	0.109

Each observed line was fitted assuming that its shape is fully described by the RGS line spread function (the intrinsic width of coronal emission lines is expected to be negligible compared to the RGS line spread function). For each wavelength window encompassing the lines to fit we took into account the continuum contribution in the best fit procedure, leaving temperature and normalization as free parameters. Given the small width of each wavelength window, the continuum component from the best fit acts as an additive constant value. This additive constant contribution, mimicking the apparent continuum level, ensures that the contributions of unresolved small lines and wings of other strong lines in the vicinity are taken into account. The measured line fluxes are listed in Table 4. In the case of spectral features suspected to be due to the blend of more than one line, we listed the different ions contributing to the observed flux.

To determine absolute abundances and EMD values, and to better constrain the hottest components of the EMD, continuum flux measurements are also needed. The continuum flux level obtained within the line fitting procedures is often an overesti-


mate of the real continuum level, because of the significant contribution of small unresolved lines, that is known to affect large portions of the RGS wavelength range. Therefore, to obtain reliable measurements of the continuum, we selected small wavelength intervals where line contamination is known to be small (on average less than 10%). The continuum flux measurements in these intervals were obtained by integrating the observed net counts. The total emissivity function associated to these continuum measurements was anyhow computed taking into account also the small fraction of emission line contribution in addition to the continuum one. The selected intervals and the corresponding measured fluxes are listed in Table 4.

We derived the EMD and the abundances corresponding to the quiescent and flaring phases by applying the Markov chain Monte Carlo method of Kashyap & Drake (1998). Flux measurements used for the EMD reconstruction are listed in Table 4. We discarded density sensitive lines and lines whose fluxes are highly uncertain because of strong blending. Measured fluxes were first converted into unabsorbed fluxes assuming a hydrogen

Table 4. Measured fluxes of emission lines and continuum intervals in the X-ray spectrum of DS Tuc.

λ^a	Ion	$\log T_{\max}^b$	quiescent		flaring		EMD ^e
			F_{obs}^c	$(F_{\text{obs}} - F_{\text{pred}}^d)/\sigma$	F_{obs}^c	$(F_{\text{obs}} - F_{\text{pred}}^d)/\sigma$	
<i>Lines</i>							
8.42	MgXII MgXII	7.00	5.7 ± 3.5	-1.3	32.4 ± 10.1	-0.3	*
9.17	MgXI	6.80	20.0 ± 3.9	2.6	32.0 ± 9.1	1.7	*
9.23	MgXI	6.80	3.2 ± 3.2	0.6	27.1 ± 8.8	2.8	*
9.31	MgXI	6.80	14.2 ± 3.4	2.9	7.7 ± 7.8	0.1	*
10.62	FeXIX FeXIX FeXVII FeXXIV FeXXIV	7.30	1.6 ± 2.6	-1.4	51.9 ± 12.8	-0.0	*
10.98	FeXXIII NeIX NaX FeXXIII FeXVII FeXXIV	7.20	22.8 ± 4.5	3.0	62.1 ± 13.8	0.4	*
11.43	FeXXII FeXXIV FeXVIII	7.20	15.3 ± 4.8	2.2	41.7 ± 13.1	1.1	*
11.74	FeXXIII	7.20	0.8 ± 2.7	-2.0	27.1 ± 12.6	-0.7	*
11.77	FeXXII	7.10	13.3 ± 4.5	1.1	20.8 ± 12.5	-0.1	*
12.13	FeXVII NeX NeX	6.75	106.4 ± 8.4	1.5	201.8 ± 18.6	-0.2	*
12.28	FeXXI FeXVII	7.05	20.9 ± 6.0	-1.3	54.0 ± 14.8	-0.0	*
12.83	FeXX FeXX FeXX	7.05	27.8 ± 5.5	1.2	41.6 ± 12.8	0.2	*
13.45	FeXIX NeIX FeXIX FeXXI	6.60	57.1 ± 7.2	0.2	127.2 ± 15.6	0.5	*
13.52	NeIX FeXIX	6.95	30.3 ± 6.5	1.5	34.9 ± 13.4	-0.4	*
13.70	NeIX	6.55	30.5 ± 6.2	1.7	48.0 ± 12.6	-0.0	*
13.74	FeXIX	6.95	13.1 ± 4.7	2.3	5.9 ± 7.6	0.2	*
13.82	FeXIX FeXVII	6.90	12.3 ± 4.0	-0.3	35.8 ± 9.2	1.5	*
14.20	FeXVIII FeXVIII FeXVIII	6.90	42.4 ± 4.8	-0.4	65.0 ± 9.7	-0.3	*
14.37	FeXVIII FeXVIII FeXVIII	6.90	15.0 ± 3.8	0.0	28.0 ± 8.7	0.6	*
14.54	FeXVIII FeXVIII FeXVIII	6.90	5.6 ± 3.5	-2.1	23.8 ± 8.6	0.5	*
15.01	FeXVII	6.75	97.9 ± 5.9	0.9	152.9 ± 11.8	0.2	*
15.18	FeXIX OVIII OVIII	6.50	19.7 ± 4.9	1.4	42.5 ± 10.6	1.2	*
15.26	FeXVII	6.75	26.2 ± 4.7	-0.1	8.2 ± 8.7	-4.0	*
16.01	FeXVIII OVIII OVIII	6.50	34.1 ± 4.5	-1.6	100.0 ± 10.4	0.9	*
16.07	FeXVIII	6.85	35.7 ± 4.5	3.8	40.0 ± 9.0	1.4	*
16.78	FeXVII	6.75	49.5 ± 4.6	-1.4	109.1 ± 10.3	1.4	*
17.05	FeXVII	6.75	76.3 ± 6.6	0.7	215.8 ± 15.7	6.0	*
17.10	FeXVII	6.70	59.9 ± 7.7	0.3	34.5 ± 14.0	-4.6	*
17.62	FeXVIII	6.90	7.9 ± 3.0	-1.5	8.7 ± 6.5	-1.6	*
18.63	OVII	6.35	9.8 ± 3.1	1.4	17.2 ± 7.4	1.0	*
18.97	OVIII OVIII	6.50	213.9 ± 8.8	0.2	516.6 ± 18.8	-0.1	*
21.60	OVII	6.30	46.4 ± 7.1	0.5	74.1 ± 14.0	0.2	*
21.81	OVII	6.30	0.7 ± 3.4	-1.8	14.2 ± 11.5	0.4	*
22.10	OVII	6.30	37.8 ± 6.8	2.5	28.8 ± 11.5	-0.1	*
24.78	NVII NVII	6.30	25.4 ± 5.3	-0.8	72.1 ± 12.0	0.5	*
33.73	CVI CVI	6.15	33.1 ± 8.9	-0.8	98.4 ± 19.2	0.5	*
<i>Continuum</i>							
[8.49÷8.90]		7.75	25.5 ± 4.2	0.7	192.1 ± 11.6	-0.0	*
[29.70÷30.36]		6.55	27.9 ± 11.4	-0.3	165.3 ± 21.3	0.1	*

^a Wavelengths (Å). ^b Temperature (K) of maximum emissivity. ^c Observed fluxes (10^{-6} ph s $^{-1}$ cm $^{-2}$) with uncertainties at the 68% confidence level. ^d Predicted fluxes considered here those computed in the low density limit. ^e Flux measurements selected for the EMD reconstruction.


Fig. 6. Emissivity functions of the ion lines (colored curves) and continuum (black curves) measured in RGS spectra.

phase, where, due to the higher line fluxes, they are determined with better precision. The EMD and the abundances derived for the X-ray emitting plasma of DS Tuc A are reported in Table 5. The two EMDs are also plotted in Fig. 7.

It is possible to constrain the X-ray emitting plasma density n_e by inspecting the diagnostics provided by He-like line

triplets and other density-sensitive line ratios. Among the emission lines detected in the flaring and quiescent RGS spectra, density-sensitive line ratios are provided by the He-like triplets of Mg XI (resonance r at 9.17 Å, intercombination i at 9.23 Å, forbidden f at 9.31 Å), Ne IX ($\lambda_r = 13.45$ Å, $\lambda_i = 13.55$ Å, $\lambda_f = 13.70$ Å), and O VII ($\lambda_r = 21.60$ Å, $\lambda_i = 21.81$ Å, and $\lambda_f = 22.10$ Å), and by the pair of Fe XVII lines at 17.05 Å and 17.10 Å (Mauche et al. 2001).

However, we did not analyze the He-like Ne IX triplet at ~ 13.5 Å, because of its severe blending with strong Fe lines makes its measured fluxes highly uncertain. The O VII triplet, that has a maximum formation temperature of ~ 2 MK, allows us to measure the density of relatively cold coronal plasma. The Mg XI and Fe XVII density diagnostics instead probe hotter plasma components, since their line maximum formation temperatures are at ~ 6 MK.

The f/i line ratio of the O VII triplet is high and compatible with the low density limits for both the quiescent and flaring phases. Density sensitive lines of Mg XI and Fe XVII, of the quiescent and flaring phases, are displayed in Fig. 8. Both the quiescent and flaring spectra in the Mg XI region are quite noisy due to the low S/N ratio of the lines, and to the significant continuum emission in this region, especially during flares. The Fe XVII lines are well exposed, with a good S/N ratio in both the quiescent and flare spectra. However, the Fe XVII lines are separated


Table 5. EMD and abundances of DS Tuc.

Fig. 7. EMDs derived from the RGS line fluxes for the quiescent (blue) and flaring phases (red). Asterisks and solid points mark the EM derived from EPIC detectors for the quiescent (blue symbol, 3T APEC) and flaring intervals, (red symbols and interval numbers, 1T APEC).

by 0.05 \AA which is comparable with the RGS spectral resolution (0.06 \AA) making the two lines only marginally resolved. Because of all these sources of uncertainty, the Mg XI and Fe XVII line fluxes were obtained by freezing line positions, and leaving the line fluxes as free parameters for the best fit of the line profiles.

The observed f/i ratio of Mg XI and the $F(17.10)/F(17.05)$ ratio of Fe XVII change significantly during the quiescent and flaring phases. During quiescent emission both Mg XI and Fe XVII have both ratios compatible with the low density limit: $f/i > 2$ and $F(17.10)/F(17.05) = 0.87^{+0.13}_{-0.12}$, both indicating $n_e < 10^{13} \text{ cm}^{-3}$. Conversely, during the flaring phases, both ratios decrease significantly ($f/i = 0.28^{+0.36}_{-0.28}$ and $F(17.10)/F(17.05) = 0.16 \pm 0.07$) indicating a high plasma density ($n_e \sim 10^{14} \text{ cm}^{-3}$), as shown in Fig. 9. This value is derived from relatively cold lines and they probe the cooler plasma in the loop which is toward its feet (Argiroffi et al. 2019). Conversely, the plasma density inferred from the emission measure and the loop volume described in Sect. 3.2 is valid for hotter plasma towards the apex of the loop during the flare peak. Also, in deriving the electron density from the flare emission measure, the choice of the volume has some degree of arbitrariness.

The abundances of elements inferred from the line fluxes, scaled to the values of the solar photosphere as a function of First Ionization Potential (FIP), are reported in Fig. 10. Apart from the high uncertainty of the abundance of C, a trend with FIP is visible with Fe, Mg and perhaps O being under abundant owing to their low FIP. The trend is in agreement with the inverse low FIP effect observed in active stars.

4. Discussion and conclusions

In this paper we have presented the analysis of an *XMM-Newton* observation of DS Tuc A which is one of the youngest known planet hosts. DS Tuc Ab is an inflated planet with a radius of

	quiescent	flaring
log T (K)	log EMD (cm^{-3})	log EMD (cm^{-3})
6.00	$48.36^{+1.50}_{-0.59}$	$49.08^{+0.89}_{-0.96}$
6.10	$49.43^{+1.05}_{-0.45}$	$49.36^{+0.96}_{-0.59}$
6.20	$50.11^{+0.77}_{-0.46}$	$50.48^{+0.45}_{-0.61}$
6.30	$50.20^{+0.59}_{-0.37}$	$50.42^{+0.58}_{-0.45}$
6.40	$51.66^{+0.99}_{-0.37}$	$51.00^{+0.38}_{-0.41}$
6.50	$51.05^{+0.47}_{-0.18}$	$51.42^{+0.37}_{-0.37}$
6.60	$51.51^{+0.45}_{-0.14}$	$52.60^{+0.06}_{-0.38}$
6.70	$52.17^{+0.16}_{-0.17}$	$52.00^{+0.48}_{-0.49}$
6.80	$52.28^{+0.32}_{-0.34}$	$52.28^{+0.49}_{-0.42}$
6.90	$51.68^{+0.50}_{-0.18}$	$52.02^{+0.37}_{-0.17}$
7.00	$52.30^{+0.11}_{-0.39}$	$52.61^{+0.09}_{-0.43}$
7.10	$52.27^{+0.15}_{-0.22}$	$52.39^{+0.33}_{-0.26}$
7.20	$51.85^{+0.22}_{-0.50}$	$52.54^{+0.26}_{-0.52}$
7.30	$50.95^{+0.37}_{-0.50}$	$52.53^{+0.27}_{-0.25}$
7.40	$50.60^{+0.31}_{-0.61}$	$52.67^{+0.31}_{-0.66}$
7.50	$49.35^{+1.23}_{-0.45}$	$53.25^{+0.67}_{-0.67}$
7.60	$49.73^{+0.83}_{-0.81}$	$52.47^{+0.40}_{-0.24}$
7.70	$48.42^{+1.80}_{-0.43}$	$52.28^{+0.34}_{-0.21}$
7.80	$47.43^{+2.57}_{-0.52}$	$51.98^{+0.69}_{-0.69}$
7.90	$47.33^{+2.80}_{-0.54}$	$52.42^{+0.45}_{-0.98}$
8.00	$47.47^{+2.47}_{-0.79}$	$51.19^{+0.89}_{-0.68}$
Elem	$A_X/A_{X\odot}$	
C	$0.73^{+0.28}_{-0.16}$	
N	$0.85^{+0.35}_{-0.15}$	
O	$0.54^{+0.16}_{-0.08}$	
Ne	$0.91^{+0.30}_{-0.16}$	
Mg	$0.41^{+0.16}_{-0.10}$	
Fe	$0.26^{+0.07}_{-0.05}$	

Abundances are in solar units (Anders & Grevesse 1989, GeCoA, 53, 97).

about $0.5 R_J$ and with mass less than $14.4 M_{\oplus}$ that will likely shrink to a size of about $2 R_{\oplus}$ in the next hundreds of Myr (Bennati et al. 2021). Observations in X-rays are important to understand the processes that can modify both dynamics and chemistry of the primary atmospheres of newly born planets. In this context, the present work allows a deep characterization of the coronal emission and time variability of the star and its effects on its close-in planet.

We detected two bright flares in DS Tuc A that permitted a time-resolved spectroscopy of the two events. The energy released during the first and second flare are about 8×10^{34} erg and 5×10^{34} erg.

Colombo et al. (2022) used TESS light curves of DS Tuc to study the rate of flares of DS Tuc in the optical band with an analysis of all available sectors through an iterative method based on Gaussian processes. They found that the frequency of flares with energy (in optical band) larger than 2×10^{32} erg amounts to about 2 per day. By using the relationship between flare energy released in the optical band and in X-rays inferred by Flaccomio et al. (2018) from the flares observed in the NGC 2264 young star forming region, this limit corresponds to X-ray energies of order of 10^{31} erg. The flares we detected in the present *XMM-Newton* observation seem very rare events that likely released energies in the optical band in excess of 5×10^{35} erg which are on the very high energy tail of the flare energies observed with TESS by Colombo et al (2022). Furthermore, in the optical band there is a significant fraction of pairs of flares separated by a few ks, similar to the separation of the two flares we observed




Fig. 8. Observed (histogram with error bars) and best fit (solid black curve) spectra, of the quiescent and flaring phases, in the regions of the Mg XI triplet and Fe XVII density sensitive lines. Best fit functions corresponding to individual line are also shown with solid grey lines. *Left panels*; *r*, *f*, and *i* labels indicate the resonance, inter-combination, and forbidden lines of the Mg XI triplet.




Fig. 9. Predicted line ratios (f/i for O VII and Mg XI and $F(17.10)/F(17.05)$ for Fe XVII) computed at different plasma densities, together with the observed line ratios observed during the flaring phase.

in X-rays. The cause could be a triggering mechanism of the same active region after the first flare has been ignited in a scenario where the corona is densely packed with magnetic structures somehow interacting with each other. The UV energy of the flares was calculated from the count rates and the conversion factor for the UVM2 filter (assuming a band width of 100 Å) in the case of a WD at 10,000 K³. These energies amount to 2.7 and 0.9×10^{33} erg for flares 1 and 2, respectively, and result a factor 30 to 50 lower than the energies released in X-rays.

In pre Main Sequence stars of Orion, X-ray flares that release energies $\geq 5.5 \times 10^{35}$ erg constitute the upper 50% of the sample of the bright flares observed in COUP (cf. Fig. 9 in Wolk et al. 2005). It is plausible that the frequency of such energetic events decreases between the age of young pre-Main Sequence stars in Orion (3 – 5 Myr) and the age of DS Tuc, about 40 Myr.

With an age of 40 Myr the X-ray activity of DS Tuc A has just started its decline in the Main Sequence phase and probed by a value of the ratio $\log L_X/L_{\text{bol}} \sim -3.55$ (Benatti et al. 2021). The average X-ray luminosity of DS Tuc A in this observation was 9.1×10^{30} erg s⁻¹ which is about a factor 10 higher than recorded

in the first *XMM-Newton* observation (Benatti et al. 2021). During the flares the ratio $\log L_X/L_{\text{bol}}$ reached the value of ≈ -2.15 or $\approx 0.7\%$ of its bolometric luminosity.

From the ephemeris of Benatti et al. (2019) we estimated that the planet was at the orbital phases ϕ between ~ -0.06 and ~ -0.11 during this observation (being $\phi = 0$ the phase of the transit and ~ -0.008 the phase of the first contact) and thus the planet was observed between 19 and 12 hours before the transit. The stellar rotation period is about 2.9 days which amounts to ≈ 250.5 ks, thus the present *XMM-Newton* exposure covered about 12% of the rotation period. The flares did not show any dimming due to self-eclipse, we infer that either they occurred in active regions near the equator and in the central portion of the stellar disk or the active regions were at very high stellar latitudes and coronal loops with sizes similar to the stellar radius anchored at high stellar latitudes could be visible even when the feet of the loops are behind the visible disk.

The values of peak luminosity are about 1.8×10^{31} erg/s, these values correspond to a flux of about $100 \text{ erg s}^{-1} \text{ cm}^{-2}$ at the surface of the planet, which is of order of 0.1 W m^{-2} . Conversely, the X-ray flux received by the planet from the quiescent star is about 0.01 W m^{-2} . An hypothetical Earth around DS Tuc A at 1 AU would have received $6.3 \times 10^{-4} \text{ W m}^{-2}$ in the band 0.3 – 10 keV, which corresponds to a GOES X6 flare in the band 1 – 8 Å. We speculate that Coronal Mass Ejections (CMEs) were associated with those flares. Drake et al. (2013) (see also Osten & Wolk 2015; Moschou et al. 2019) give an estimate of the mass and kinetic energy associated with powerful flares observed in the Sun. From their equation 1, which applies to young stellar coroneae in the saturated regime like in the case of DS Tuc A, we estimated that the mass associated with CMEs was around $5 \times 10^{-15} M_{\odot}$. We also infer that the kinetic energy associated with the CMEs is $\approx 10^{34}$ erg. From these numbers we also infer a velocity of the CME material of order of 1000 km/s. With these velocities the CME would have reached DS Tuc Ab in about 3.3 hr after the ignition of the flares. Simulations of CMEs hitting close-in planets at a few stellar radii from their stars (see Alvarado-Gómez et al. 2022; Hazra et al. 2022) demonstrate that the CMEs associated with the flares could be even more disruptive for the outer atmosphere resulting in dramatic changes of the rate of evaporation with respect to the effect of a steady stellar wind or from X-ray irradiation alone due to a flare.

The RGS spectra enable us to reconstruct the distribution of Emission Measure (EMD) for the quiescent and flaring states.

³ See <https://www.cosmos.esa.int/web/xmm-newton/sas-watchout-uvflux>.




Fig. 10. Ratio $A_X/A_{X\odot}$ vs. first ionization potential derived from the abundances analysis of RGS spectra

The EMDs are very similar in the range $6 < \log T < 7.1$ while they differ significantly above $\log T > 7.1$ with the maximum difference occurring at $\log T \approx 7.5$. The difference is due to the dense plasma heated during the flares. The integral of the difference between the flare and quiescent phase EMDs in the range of temperature $7 < \log T < 8$ amounts to $\log EM \approx 53.5$ which is consistent, on average, with the values inferred from the time resolved spectroscopy of EPIC spectra. Qualitatively, hard X-rays produced during the flares can penetrate more deeply in the planet's atmosphere along with a cascade of electrons that induce further dissociation and photo-chemistry. Such a mechanism can be non linear if the atmosphere is unable to restore its state before the next flare. Louca et al. (2022) modeled the response of several planetary atmospheres to flares determining the conditions under which the initial conditions are restored and the consequent enhancement of several species. In this respect, a more systematic X-ray monitoring of stars with close in planets would be beneficial for the comprehension of the formation and evolution of planetary atmospheres.

Acknowledgements. We thank the anonymous referee for their useful comments and suggestions that improved the quality of this manuscript. IP, AM, SC and GM acknowledge financial support from the ASI-INAF agreement n.2018-16-HH.0, and from the ARIEL ASI-INAF agreement n.2021-5-HH.0. SJW was supported by the Chandra X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of the National Aeronautics Space Administration under contract NAS8-03060. Based on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA.

References

Alvarado-Gómez, J. D., Cohen, O., Drake, J. J., et al. 2022, *ApJ*, 928, 147

Argiroffi, C., Reale, F., Drake, J. J., et al. 2019, *Nature Astronomy*, 3, 742

Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17

Benatti, S., Damasso, M., Borsa, F., et al. 2021, *A&A*, 650, A66

Benatti, S., Nardiello, D., Malavolta, L., et al. 2019, *A&A*, 630, A81

Cecchi-Pestellini, C., Ciaravella, A., Micela, G., & Penz, T. 2009, *A&A*, 496, 863

Colombo, S., Petralia, A., & Micela, G. 2022, arXiv e-prints, arXiv:2202.13615

Dennis, B. R. & Zarro, D. M. 1993, *Sol. Phys.*, 146, 177

Dere, K. P., Landi, E., Mason, H. E., Monsignori Fossi, B. C., & Young, P. R. 1997, *A&AS*, 125, 149

Drake, J. J., Cohen, O., Yashiro, S., & Gopalswamy, N. 2013, *ApJ*, 764, 170

Favata, F., Giardino, G., Micela, G., Sciortino, S., & Damiani, F. 2003, *A&A*, 403, 187

Feigelson, E. D. & Montmerle, T. 1999, *ARA&A*, 37, 363

Flaccomio, E., Micela, G., Sciortino, S., et al. 2018, *A&A*, 620, A55

Hazra, G., Vidotto, A. A., Carolan, S., Villarreal D'Angelo, C., & Manchester, W. 2022, *MNRAS*, 509, 5858

Hudson, H. S. 1991, *Sol. Phys.*, 133, 357

Kahler, S. W., Moore, R. L., Kane, S. R., & Zirin, H. 1988, *ApJ*, 328, 824

Kashyap, V. & Drake, J. J. 1998, *ApJ*, 503, 450

Kashyap, V. & Drake, J. J. 2000, *Bulletin of the Astronomical Society of India*, 28, 475

Kay, C., Airapetian, V. S., Lüftinger, T., & Kochukhov, O. 2019, *ApJ*, 886, L37

Kublyshkina, D., Fossati, L., Erkaev, N. V., et al. 2018, *ApJ*, 866, L18

Louca, A. J., Miguel, Y., Tsai, S.-M., et al. 2022, arXiv e-prints, arXiv:2204.10835

Maggio, A., Pallavicini, R., Reale, F., & Tagliaferri, G. 2000, *A&A*, 356, 627

Mauche, C. W., Liedahl, D. A., & Fournier, K. B. 2001, *ApJ*, 560, 992

Moschou, S.-P., Drake, J. J., Cohen, O., et al. 2019, *ApJ*, 877, 105

Namekata, K., Sakaue, T., Watanabe, K., et al. 2017, *ApJ*, 851, 91

Neupert, W. M. 1968, *ApJ*, 153, L59

Newton, E. R., Mann, A. W., Tofflemire, B. M., et al. 2019, *ApJ*, 880, L17

Osten, R. A. & Wolk, S. J. 2015, *ApJ*, 809, 79

Pallavicini, R., Golub, L., Rosner, R., et al. 1981, *ApJ*, 248, 279

Penz, T. & Micela, G. 2008, *A&A*, 479, 579

Pizzolato, N., Maggio, A., Micela, G., Sciortino, S., & Ventura, P. 2003, *A&A*, 397, 147

Reale, F. 2007, *A&A*, 471, 271

Reale, F. 2014, *Living Reviews in Solar Physics*, 11, 4

Vaiana, G. S., Fabbiano, G., Giacconi, R., et al. 1981, *ApJ*, 245, 163

Veronig, A., Temmer, M., Hanslmeier, A., Otruba, W., & Messerotti, M. 2002, *A&A*, 382, 1070

Wolk, S. J., Harnden, Jr., F. R., Flaccomio, E., et al. 2005, *ApJS*, 160, 423